

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
8 March 2007 (08.03.2007)

PCT

(10) International Publication Number  
**WO 2007/028129 A1**

(51) International Patent Classification:  
C07D 471/04 (2006.01) A61P 37/02 (2006.01)  
A61K 31/437 (2006.01)

(21) International Application Number:  
PCT/US2006/034427

(22) International Filing Date:  
1 September 2006 (01.09.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/713,704 2 September 2005 (02.09.2005) US

(71) Applicant (for all designated States except US): **3M INNOVATIVE PROPERTIES COMPANY** [US/US];  
3M Center, Post Office Box 33427, Saint Paul, Minnesota  
55133-3427 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DELLARIA, Joseph, F., Jr.** [US/US]; 3M Center, Post Office Box 33427, Saint Paul, Minnesota 55133-3427 (US). **MOSER, William, H.** [US/US]; 3M Center, Post Office Box 33427, Saint Paul, Minnesota 55133-3427 (US). **RADMER, Matthew, R.** [US/US]; 3M Center, Post Office Box 33427, Saint Paul, Minnesota 55133-3427 (US). **GRIES-GRABER, George, W.** [US/US]; 3M Center, Post Office Box 33427, Saint Paul, Minnesota 55133-3427 (US).

(74) Agents: **ERSFELD, Dean, A.** et al.; 3M Center, Office of Intellectual Property Counsel, Post Office Box 33427, Saint Paul, Minnesota 55133-3427 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HYDROXY SUBSTITUTED 1H-IMIDAZOPYRIDINES AND METHODS

(57) Abstract: Hydroxy substituted 1H-imidazo[4,5-c]pyridin-4-amines, with a hydroxy substituent at the 2-position, pharmaceutical compositions containing these compounds, methods of making the compounds, intermediates, and methods of use of these compounds as immunomodulators, for inducing cytokine biosynthesis in animals and in the treatment of diseases including viral and neoplastic diseases, are disclosed.

WO 2007/028129 A1

## HYDROXY SUBSTITUTED 1H-IMIDAZOPYRIDINES AND METHODS

## CROSS REFERENCE TO RELATED APPLICATIONS


5           The present invention claims priority to U.S. Provisional Application Serial No. 60/713,704, filed September 2, 2005, which is incorporated herein by reference.

## BACKGROUND


10           Certain compounds have been found to be useful as immune response modifiers (IRMs), rendering them useful in the treatment of a variety of disorders. However, there continues to be interest in and a need for compounds that have the ability to modulate the immune response, by induction of cytokine biosynthesis or other means.

## SUMMARY OF THE INVENTION


15           It has now been found that certain 2-hydroxy-1H-imidazo[4,5-c]pyridin-4-amines modulate cytokine biosynthesis. In one aspect, the present invention provides compounds, which are of the following Formulas I, II, and III:



I



II



III

wherein  $R_1$ ,  $R_A$ ,  $R_B$ ,  $G_1$ , and  $G_2$  are as defined below; and pharmaceutically acceptable salts thereof.

5 The compounds or salts of Formulas I, II, and III are useful as IRMs due to their ability to modulate cytokine biosynthesis (e.g., induce the biosynthesis or production of one or more cytokines) and otherwise modulate the immune response when administered to animals. In some embodiments, compounds or salts of Formula I can be especially useful as immune response modifiers due to their ability to selectively induce interferon ( $\alpha$ ) (IFN- $\alpha$ ), thus providing a benefit over compounds that also induce pro-inflammatory  
10 cytokines (e.g. TNF- $\alpha$ ) or that induce pro-inflammatory cytokines at higher levels. The ability to modulate cytokine biosynthesis makes the compounds useful in the treatment of a variety of conditions such as viral diseases and neoplastic diseases, that are responsive to such changes in the immune response.

In another aspect, the present invention also provides pharmaceutical compositions  
15 containing the compounds of Formulas I, II, and/or III, and methods of inducing cytokine biosynthesis in animal cells, selectively inducing IFN- $\alpha$  in animal cells, treating a viral disease in an animal, and/or treating a neoplastic disease in an animal by administering to the animal one or more compounds of the Formulas I, II, and/or III, and/or pharmaceutically acceptable salts thereof.

20 In another aspect, the invention provides methods of synthesizing the compounds of Formulas I, II, and III and intermediate compounds useful in the synthesis of these compounds.


As used herein, "a", "an", "the", "at least one", and "one or more" are used interchangeably.

25 The terms "comprising" and variations thereof do not have a limiting meaning where these terms appear in the description and claims.


The above summary of the present invention is not intended to describe each disclosed embodiment or every implementation of the present invention. The description that follows more particularly exemplifies illustrative embodiments. Guidance is also  
30 provided herein through lists of examples, which can be used in various combinations. In each instance, the recited list serves only as a representative group and should not be interpreted as an exclusive list.

# DETAILED DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS OF THE INVENTION


The present invention provides compounds of the following Formulas I, II, and III:



I




II



III

wherein  $R_1$ ,  $R_A$ ,  $R_B$ ,  $G_1$ , and  $G_2$  are as defined below; and pharmaceutically acceptable salts thereof.

In one embodiment, the present invention provides a compound of the following Formula I:



I

wherein:

$R_A$  and  $R_B$  are each independently selected from the group consisting of:

hydrogen,  
 halogen,  
 alkenyl,  
 amino,  
 5 -R<sub>11</sub>,  
 -O-R<sub>11</sub>,  
 -S-R<sub>11</sub>, and  
 -N(R<sub>9a</sub>)(R<sub>11</sub>);

R<sub>11</sub> is selected from the group consisting of alkyl, alkoxyalkylenyl,  
 10 hydroxyalkylenyl, aryl, arylalkylenyl, heteroaryl, heteroarylalkylenyl, heterocyclyl, and  
 heterocyclylalkylenyl, each of which is unsubstituted or substituted by one or more  
 substituents independently selected from the group consisting of alkyl; alkoxy; hydroxy;  
 hydroxyalkyl; aryl; aryloxy; arylalkyleneoxy; heteroaryl; heteroaryloxy;  
 heteroarylalkyleneoxy; halogen; haloalkyl; haloalkoxy; mercapto; nitro; cyano;  
 15 heterocyclyl; amino; alkylamino; dialkylamino; and, in the case of alkyl, heterocyclyl, and  
 heterocyclylalkylenyl, oxo;

R<sub>9a</sub> is selected from the group consisting of hydrogen and C<sub>1-4</sub> alkyl;

R<sub>1</sub> is selected from the group consisting of:



-R<sub>4</sub>,  
 20 -X-R<sub>4</sub>,  
 -X-Y-R<sub>4</sub>,  
 -X-Y-X-Y-R<sub>4</sub>,  
 -X-R<sub>5</sub>,  
 -N(R<sub>1</sub>')-Q-R<sub>4</sub>,  
 25 -N(R<sub>1</sub>')-X<sub>1</sub>-Y<sub>1</sub>-R<sub>4</sub>, and  
 -N(R<sub>1</sub>')-X<sub>1</sub>-R<sub>5a</sub>;

X is selected from the group consisting of alkylene, alkenylene, alkynylene,  
 arylene, heteroarylene, and heterocyclylene wherein the alkylene, alkenylene, and  
 alkynylene groups can be optionally interrupted or terminated by arylene, heteroarylene or  
 30 heterocyclylene and optionally interrupted by one or more -O- groups;


X<sub>1</sub> is C<sub>2-20</sub> alkylene;

Y is selected from the group consisting of:

-O-,  
 -S(O)<sub>0-2</sub>-,  
 -S(O)<sub>2</sub>-N(R<sub>8</sub>)-,  
 -C(R<sub>6</sub>)-,  
 5 -C(R<sub>6</sub>)-O-,  
 -O-C(R<sub>6</sub>)-,  
 -O-C(O)-O-,  
 -N(R<sub>8</sub>)-Q-,  
 -C(R<sub>6</sub>)-N(R<sub>8</sub>)-,  
 10 -O-C(R<sub>6</sub>)-N(R<sub>8</sub>)-,  
 -C(R<sub>6</sub>)-N(OR<sub>9</sub>)-,  
 -O-N(R<sub>8</sub>)-Q-,  
 -O-N=C(R<sub>4</sub>)-,  
 -C(=N-O-R<sub>8</sub>)-,  
 15 -CH(-N(-O-R<sub>8</sub>)-Q-R<sub>4</sub>)-,




Y<sub>1</sub> is selected from the group consisting of -O-, -S(O)<sub>0-2</sub>-, -S(O)<sub>2</sub>-N(R<sub>8</sub>)-,

-N(R<sub>8</sub>)-Q-, -C(R<sub>6</sub>)-N(R<sub>8</sub>)-, -O-C(R<sub>6</sub>)-N(R<sub>8</sub>)-, and  

 ;


$R_1'$  is selected from the group consisting of hydrogen,  $C_{1-20}$  alkyl, hydroxy- $C_{2-20}$  alkyl, and alkoxy- $C_{2-20}$  alkyl;

$R_4$  is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, arylalkyl, aryloxyalkyl, alkylarylenyl, heteroaryl, heteroarylalkyl, heteroaryloxyalkyl, alkylheteroarylenyl, and heterocycl, wherein the alkyl, alkenyl, alkynyl, aryl, arylalkyl, aryloxyalkyl, alkylarylenyl, heteroaryl, heteroarylalkyl, heteroaryloxyalkyl, alkylheteroarylenyl, and heterocycl groups can be unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl; alkoxy; hydroxyalkyl; haloalkyl; haloalkoxy; halogen; nitro; hydroxy; mercapto; cyano; aryl; aryloxy; arylalkyleneoxy; heteroaryl; heteroaryloxy; heteroarylalkyleneoxy; heterocycl; amino; alkylamino; dialkylamino; (dialkylamino)alkyleneoxy; and, in the case of alkyl, alkenyl, alkynyl, and heterocycl, oxo;

$R_5$  is selected from the group consisting of:



$R_{5a}$  is selected from the group consisting of:



$R_6$  is selected from the group consisting of =O and =S;

$R_7$  is  $C_{2-7}$  alkylene;

$R_8$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl, hydroxy- $C_{1-10}$  alkyl,  $C_{1-10}$  alkoxy- $C_{1-10}$  alkyl, aryl- $C_{1-10}$  alkyl, and heteroaryl- $C_{1-10}$  alkyl;

$R_9$  is selected from the group consisting of hydrogen and alkyl;

$R_{10}$  is  $C_{3-8}$  alkylene;

A is selected from the group consisting of  $-\text{CH}_2-$ ,  $-\text{O}-$ ,  $-\text{C(O)}-$ ,  $-\text{S(O)}_{0-2}-$ , and

-N(-Q-R<sub>4</sub>)-;

A' is selected from the group consisting of -O-, -S(O)<sub>0-2</sub>-, -N(-Q-R<sub>4</sub>)-, and -CH<sub>2</sub>-;


Q is selected from the group consisting of a bond, -C(R<sub>6</sub>)-, -C(R<sub>6</sub>)-C(R<sub>6</sub>)-, -S(O)<sub>2</sub>-, -C(R<sub>6</sub>)-N(R<sub>8</sub>)-W-, -S(O)<sub>2</sub>-N(R<sub>8</sub>)-, -C(R<sub>6</sub>)-O-, -C(R<sub>6</sub>)-S-, and -C(R<sub>6</sub>)-N(OR<sub>9</sub>)-;

5 V is selected from the group consisting of -C(R<sub>6</sub>)-, -O-C(R<sub>6</sub>)-, -N(R<sub>8</sub>)-C(R<sub>6</sub>)-, and -S(O)<sub>2</sub>-;

W is selected from the group consisting of a bond, -C(O)-, and -S(O)<sub>2</sub>-; and

a and b are independently integers from 1 to 6 with the proviso that a + b is ≤ 7; or a pharmaceutically acceptable salt thereof.

10 In one embodiment, the present invention provides a compound of the following Formula II, which is a prodrug:



II

wherein:

15 G<sub>1</sub> is selected from the group consisting of:

-C(O)-R',  
 α-aminoacyl,  
 α-aminoacyl-α-aminoacyl,  
 -C(O)-O-R',  
 20 -C(O)-N(R'')R',  
 -C(=NY')-R',  
 -CH(OH)-C(O)-OY',  
 -CH(OC<sub>1-4</sub> alkyl)Y<sub>0</sub>,  
 -CH<sub>2</sub>Y<sub>2</sub>, and  
 25 -CH(CH<sub>3</sub>)Y<sub>2</sub>;

R' and R'' are independently selected from the group consisting of C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, phenyl, and benzyl, each of which may be unsubstituted or substituted by one or more substituents independently selected from the group consisting of halogen, hydroxy, nitro, cyano, carboxy, C<sub>1-6</sub> alkyl, C<sub>1-4</sub> alkoxy, aryl, heteroaryl, aryl-C<sub>1-4</sub> alkylenyl,



heteroaryl-C<sub>1-4</sub> alkylenyl, halo-C<sub>1-4</sub> alkylenyl, halo-C<sub>1-4</sub> alkoxy, -O-C(O)-CH<sub>3</sub>,  
 -C(O)-O-CH<sub>3</sub>, -C(O)-NH<sub>2</sub>, -O-CH<sub>2</sub>-C(O)-NH<sub>2</sub>, -NH<sub>2</sub>, and -S(O)<sub>2</sub>-NH<sub>2</sub>, with the proviso  
 that R'' can also be hydrogen;

$\alpha$ -aminoacyl is an  $\alpha$ -aminoacyl group derived from an amino acid selected from  
 5 the group consisting of racemic, D-, and L-amino acids;

Y' is selected from the group consisting of hydrogen, C<sub>1-6</sub> alkyl, and benzyl;

Y<sub>0</sub> is selected from the group consisting of C<sub>1-6</sub> alkyl, carboxy-C<sub>1-6</sub> alkylenyl,  
 amino-C<sub>1-4</sub> alkylenyl, mono-*N*-C<sub>1-6</sub> alkylamino-C<sub>1-4</sub> alkylenyl, and  
 di-*N,N*-C<sub>1-6</sub> alkylamino-C<sub>1-4</sub> alkylenyl;

10 Y<sub>2</sub> is selected from the group consisting of mono-*N*-C<sub>1-6</sub> alkylamino,  
 di-*N,N*-C<sub>1-6</sub> alkylamino, morpholin-4-yl, piperidin-1-yl, pyrrolidin-1-yl, and  
 4-C<sub>1-4</sub> alkylpiperazin-1-yl;

R<sub>A</sub> and R<sub>B</sub> are each independently selected from the group consisting of:

hydrogen,  
 15 halogen,  
 alkenyl,  
 amino,  
 -R<sub>11</sub>,  
 -O-R<sub>11</sub>,  
 20 -S-R<sub>11</sub>, and  
 -N(R<sub>9a</sub>)(R<sub>11</sub>);

R<sub>11</sub> is selected from the group consisting of alkyl, alkoxyalkylenyl,  
 hydroxyalkylenyl, aryl, arylalkylenyl, heteroaryl, heteroarylalkylenyl, heterocyclyl, and  
 heterocyclylalkylenyl, each of which is unsubstituted or substituted by one or more  
 25 substituents independently selected from the group consisting of alkyl; alkoxy; hydroxy;  
 hydroxyalkyl; aryl; aryloxy; arylalkyleneoxy; heteroaryl; heteroaryloxy;  
 heteroarylalkyleneoxy; halogen; haloalkyl; haloalkoxy; mercapto; nitro; cyano;  
 heterocyclyl; amino; alkylamino; dialkylamino; and, in the case of alkyl, heterocyclyl, and  
 heterocyclylalkylenyl, oxo;

30 R<sub>9a</sub> is selected from the group consisting of hydrogen and C<sub>1-4</sub> alkyl;

R<sub>1</sub> is selected from the group consisting of:

-R<sub>4</sub>,


$-X-R_4$ ,  
 $-X-Y-R_4$ ,  
 $-X-Y-X-Y-R_4$ ,  
 $-X-R_5$ ,  
 $-N(R_1')-Q-R_4$ ,  
 $-N(R_1')-X_1-Y_1-R_4$ , and  
 $-N(R_1')-X_1-R_{5a}$ ;


X is selected from the group consisting of alkylene, alkenylene, alkynylene,  
 arylene, heteroarylene, and heterocyclylene wherein the alkylene, alkenylene, and  
 alkynylene groups can be optionally interrupted or terminated by arylene, heteroarylene or  
 heterocyclylene and optionally interrupted by one or more -O- groups;

$X_1$  is  $C_{2-20}$  alkylene;


Y is selected from the group consisting of:

$-O-$ ,  
 $-S(O)_{0-2}-$ ,  
 $-S(O)_2-N(R_8)-$ ,  
 $-C(R_6)-$ ,  
 $-C(R_6)-O-$ ,  
 $-O-C(R_6)-$ ,  
 $-O-C(O)-O-$ ,  
 $-N(R_8)-Q-$ ,  
 $-C(R_6)-N(R_8)-$ ,  
 $-O-C(R_6)-N(R_8)-$ ,  
 $-C(R_6)-N(OR_9)-$ ,  
 $-O-N(R_8)-Q-$ ,  
 $-O-N=C(R_4)-$ ,  
 $-C(=N-O-R_8)-$ ,  
 $-CH(-N(-O-R_8)-Q-R_4)-$ ,





5  $Y_1$  is selected from the group consisting of -O-, -S(O)<sub>0-2</sub>-, -S(O)<sub>2</sub>-N(R<sub>8</sub>)-,




$R_1'$  is selected from the group consisting of hydrogen, C<sub>1-20</sub> alkyl, hydroxy-C<sub>2-20</sub> alkylenyl, and alkoxy-C<sub>2-20</sub> alkylenyl;


10  $R_4$  is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl, heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl wherein the alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl, heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl groups can be unsubstituted or substituted by one or more substituents independently selected

15 from the group consisting of alkyl; alkoxy; hydroxyalkyl; haloalkyl; haloalkoxy; halogen; nitro; hydroxy; mercapto; cyano; aryl; aryloxy; arylalkyleneoxy; heteroaryl; heteroaryloxy; heteroarylalkyleneoxy; heterocyclyl; amino; alkylamino; dialkylamino; (dialkylamino)alkyleneoxy; and, in the case of alkyl, alkenyl, alkynyl, and heterocyclyl, oxo;

20  $R_5$  is selected from the group consisting of:



R<sub>5a</sub> is selected from the group consisting of:



5 R<sub>6</sub> is selected from the group consisting of =O and =S;

R<sub>7</sub> is C<sub>2-7</sub> alkylene;

R<sub>8</sub> is selected from the group consisting of hydrogen, C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, hydroxy-C<sub>1-10</sub> alkylenyl, C<sub>1-10</sub> alkoxy-C<sub>1-10</sub> alkylenyl, aryl-C<sub>1-10</sub> alkylenyl, and heteroaryl-C<sub>1-10</sub> alkylenyl;

10 R<sub>9</sub> is selected from the group consisting of hydrogen and alkyl;

R<sub>10</sub> is C<sub>3-8</sub> alkylene;

A is selected from the group consisting of -CH<sub>2</sub>-, -O-, -C(O)-, -S(O)<sub>0-2</sub>-, and -N(-Q-R<sub>4</sub>)-;

A' is selected from the group consisting of -O-, -S(O)<sub>0-2</sub>-, -N(-Q-R<sub>4</sub>)-, and -CH<sub>2</sub>-;


15 Q is selected from the group consisting of a bond, -C(R<sub>6</sub>)-, -C(R<sub>6</sub>)-C(R<sub>6</sub>)-, -S(O)<sub>2</sub>-, -C(R<sub>6</sub>)-N(R<sub>8</sub>)-W-, -S(O)<sub>2</sub>-N(R<sub>8</sub>)-, -C(R<sub>6</sub>)-O-, -C(R<sub>6</sub>)-S-, and -C(R<sub>6</sub>)-N(OR<sub>9</sub>)-;

V is selected from the group consisting of -C(R<sub>6</sub>)-, -O-C(R<sub>6</sub>)-, -N(R<sub>8</sub>)-C(R<sub>6</sub>)-, and -S(O)<sub>2</sub>-;

W is selected from the group consisting of a bond, -C(O)-, and -S(O)<sub>2</sub>-; and

20 a and b are independently integers from 1 to 6 with the proviso that a + b is ≤ 7; or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention provides a compound of the following Formula III:



III

wherein:

G<sub>2</sub> is selected from the group consisting of:

- 5                   -X<sub>2</sub>-C(O)-R',  
                   α-aminoacyl,  
                   α-aminoacyl-α-aminoacyl,  
                   -X<sub>2</sub>-C(O)-O-R',  
                   -C(O)-N(R'')R', and  
 10                  -S(O)<sub>2</sub>-R';

X<sub>2</sub> is selected from the group consisting of a bond; -CH<sub>2</sub>-O-; -CH(CH<sub>3</sub>)-O-;  
 -C(CH<sub>3</sub>)<sub>2</sub>-O-; and, in the case of -X<sub>2</sub>-C(O)-O-R', -CH<sub>2</sub>-NH-;

R' and R'' are independently selected from the group consisting of C<sub>1-10</sub> alkyl,  
 C<sub>3-7</sub> cycloalkyl, phenyl, and benzyl, each of which may be unsubstituted or substituted by  
 15 one or more substituents independently selected from the group consisting of halogen,  
 hydroxy, nitro, cyano, carboxy, C<sub>1-6</sub> alkyl, C<sub>1-4</sub> alkoxy, aryl, heteroaryl, aryl-C<sub>1-4</sub> alkylenyl,  
 heteroaryl-C<sub>1-4</sub> alkylenyl, halo-C<sub>1-4</sub> alkylenyl, halo-C<sub>1-4</sub> alkoxy, -O-C(O)-CH<sub>3</sub>,  
 -C(O)-O-CH<sub>3</sub>, -C(O)-NH<sub>2</sub>, -O-CH<sub>2</sub>-C(O)-NH<sub>2</sub>, -NH<sub>2</sub>, and -S(O)<sub>2</sub>-NH<sub>2</sub>, with the proviso  
 that R'' can also be hydrogen;

20                  α-aminoacyl is an α-aminoacyl group derived from an amino acid selected from  
 the group consisting of racemic, D-, and L-amino acids;

R<sub>A</sub> and R<sub>B</sub> are each independently selected from the group consisting of:

- hydrogen,  
                   halogen,  
 25                  alkenyl,  
                   amino,  
                   -R<sub>11</sub>,  
                   -O-R<sub>11</sub>,  
                   -S-R<sub>11</sub>, and

-N(R<sub>9a</sub>)(R<sub>11</sub>);

R<sub>11</sub> is selected from the group consisting of alkyl, alkoxyalkylenyl, hydroxyalkylenyl, aryl, arylalkylenyl, heteroaryl, heteroarylalkylenyl, heterocyclyl, and heterocyclylalkylenyl, each of which is unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl; alkoxy; hydroxy; hydroxyalkyl; aryl; aryloxy; arylalkyleneoxy; heteroaryl; heteroaryloxy; heteroarylalkyleneoxy; halogen; haloalkyl; haloalkoxy; mercapto; nitro; cyano; heterocyclyl; amino; alkylamino; dialkylamino; and, in the case of alkyl, heterocyclyl, and heterocyclylalkylenyl, oxo;

10 R<sub>9a</sub> is selected from the group consisting of hydrogen and C<sub>1-4</sub> alkyl;

R<sub>1</sub> is selected from the group consisting of:

-R<sub>4</sub>,

-X-R<sub>4</sub>,

-X-Y-R<sub>4</sub>,

15 -X-Y-X-Y-R<sub>4</sub>,

-X-R<sub>5</sub>,

-N(R<sub>1</sub>')-Q-R<sub>4</sub>,

-N(R<sub>1</sub>')-X<sub>1</sub>-Y<sub>1</sub>-R<sub>4</sub>, and

-N(R<sub>1</sub>')-X<sub>1</sub>-R<sub>5a</sub>;

20 X is selected from the group consisting of alkylene, alkenylene, alkynylene, arylene, heteroarylene, and heterocyclylene wherein the alkylene, alkenylene, and alkynylene groups can be optionally interrupted or terminated by arylene, heteroarylene or heterocyclylene and optionally interrupted by one or more -O- groups;

X<sub>1</sub> is C<sub>2-20</sub> alkylene;

25 Y is selected from the group consisting of:

-O-,

-S(O)<sub>0-2</sub>-,







-S(O)<sub>2</sub>-N(R<sub>8</sub>)-,

-C(R<sub>6</sub>)-,


30 -C(R<sub>6</sub>)-O-,

-O-C(R<sub>6</sub>)-,

-O-C(O)-O-,



$Y_1$  is selected from the group consisting of  $-O-$ ,  $-S(O)_{0-2}-$ ,  $-S(O)_2-N(R_8)-$ ,




$-N(R_8)-Q-$ ,  $-C(R_6)-N(R_8)-$ ,  $-O-C(R_6)-N(R_8)-$ , and

$R_1'$  is selected from the group consisting of hydrogen,  $C_{1-20}$  alkyl, hydroxy- $C_{2-20}$  alkyl, and alkoxy- $C_{2-20}$  alkyl;


$R_4$  is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl, heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl wherein the alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl, heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl groups

can be unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl; alkoxy; hydroxyalkyl; haloalkyl; haloalkoxy; halogen; nitro; hydroxy; mercapto; cyano; aryl; aryloxy; arylalkyleneoxy; heteroaryl; heteroaryloxy; heteroarylalkyleneoxy; heterocyclyl; amino; alkylamino; dialkylamino; (dialkylamino)alkyleneoxy; and, in the case of alkyl, alkenyl, alkynyl, and heterocyclyl, oxo;

R<sub>5</sub> is selected from the group consisting of:



10 R<sub>5a</sub> is selected from the group consisting of:



R<sub>6</sub> is selected from the group consisting of =O and =S;

R<sub>7</sub> is C<sub>2-7</sub> alkylene;

15 R<sub>8</sub> is selected from the group consisting of hydrogen, C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, hydroxy-C<sub>1-10</sub> alkylenyl, C<sub>1-10</sub> alkoxy-C<sub>1-10</sub> alkylenyl, aryl-C<sub>1-10</sub> alkylenyl, and heteroaryl-C<sub>1-10</sub> alkylenyl;

R<sub>9</sub> is selected from the group consisting of hydrogen and alkyl;

R<sub>10</sub> is C<sub>3-8</sub> alkylene;

20 A is selected from the group consisting of -CH<sub>2</sub>-, -O-, -C(O)-, -S(O)<sub>0-2</sub>-, and -N(-Q-R<sub>4</sub>)-;

A' is selected from the group consisting of -O-, -S(O)<sub>0-2</sub>-, -N(-Q-R<sub>4</sub>)-, and -CH<sub>2</sub>-;

Q is selected from the group consisting of a bond, -C(R<sub>6</sub>)-, -C(R<sub>6</sub>)-C(R<sub>6</sub>)-, -S(O)<sub>2</sub>-, -C(R<sub>6</sub>)-N(R<sub>8</sub>)-W-, -S(O)<sub>2</sub>-N(R<sub>8</sub>)-, -C(R<sub>6</sub>)-O-, -C(R<sub>6</sub>)-S-, and -C(R<sub>6</sub>)-N(OR<sub>9</sub>)-;

25 V is selected from the group consisting of -C(R<sub>6</sub>)-, -O-C(R<sub>6</sub>)-, -N(R<sub>8</sub>)-C(R<sub>6</sub>)-, and -S(O)<sub>2</sub>-;

W is selected from the group consisting of a bond, -C(O)-, and -S(O)<sub>2</sub>-; and



a and b are independently integers from 1 to 6 with the proviso that  $a + b \leq 7$ ; or a pharmaceutically acceptable salt thereof.

In some embodiments, compounds of Formula III are prodrugs.

For any of the compounds presented herein, each one of the following variables (e.g.,  $R_1$ ,  $R_A$ ,  $G_1$ ,  $G_2$ ,  $R_4$ ,  $R_{11}$ ,  $X$ ,  $X_1$ ,  $Y$ ,  $Y_1$ ,  $A$ ,  $Q$ , and so on) in any of its embodiments can be combined with any one or more of the other variables in any of their embodiments and associated with any one of the formulas described herein, as would be understood by one of skill in the art. Each of the resulting combinations of variables is an embodiment of the present invention.

For certain embodiments, e.g., of Formula II,  $G_1$  is selected from the group consisting of  $-C(O)-R'$ ,  $\alpha$ -aminoacyl,  $\alpha$ -aminoacyl- $\alpha$ -aminoacyl,  $-C(O)-O-R'$ ,  $-C(O)-N(R'')R'$ ,  $-C(=NY')-R'$ ,  $-CH(OH)-C(O)-OY'$ ,  $-CH(OC_{1-4} \text{ alkyl})Y_0$ ,  $-CH_2Y_2$ , and  $-CH(CH_3)Y_2$ . For certain of these embodiments,  $R'$  and  $R''$  are independently selected from the group consisting of  $C_{1-10}$  alkyl,  $C_{3-7}$  cycloalkyl, phenyl, and benzyl, each of which may be unsubstituted or substituted by one or more substituents independently selected from the group consisting of halogen, hydroxy, nitro, cyano, carboxy,  $C_{1-6}$  alkyl,  $C_{1-4}$  alkoxy, aryl, heteroaryl, aryl- $C_{1-4}$  alkylenyl, heteroaryl- $C_{1-4}$  alkylenyl, halo- $C_{1-4}$  alkylenyl, halo- $C_{1-4}$  alkoxy,  $-O-C(O)-CH_3$ ,  $-C(O)-O-CH_3$ ,  $-C(O)-NH_2$ ,  $-O-CH_2-C(O)-NH_2$ ,  $-NH_2$ , and  $-S(O)_2-NH_2$ , with the proviso that  $R''$  can also be hydrogen;

$\alpha$ -aminoacyl is an  $\alpha$ -aminoacyl group derived from an amino acid selected from the group consisting of racemic, D-, and L-amino acids;

$Y'$  is selected from the group consisting of hydrogen,  $C_{1-6}$  alkyl, and benzyl;

$Y_0$  is selected from the group consisting of  $C_{1-6}$  alkyl, carboxy- $C_{1-6}$  alkylenyl, amino- $C_{1-4}$  alkylenyl, mono- $N$ - $C_{1-6}$  alkylamino- $C_{1-4}$  alkylenyl, and di- $N,N$ - $C_{1-6}$  alkylamino- $C_{1-4}$  alkylenyl; and

$Y_2$  is selected from the group consisting of mono- $N$ - $C_{1-6}$  alkylamino, di- $N,N$ - $C_{1-6}$  alkylamino, morpholin-4-yl, piperidin-1-yl, pyrrolidin-1-yl, and 4- $C_{1-4}$  alkylpiperazin-1-yl.

For certain embodiments, including any one of the above embodiments of Formula II,  $G_1$  is selected from the group consisting of  $-C(O)-R'$ ,  $\alpha$ -aminoacyl, and  $-C(O)-O-R'$ .

For certain embodiments, including any one of the above embodiments of Formula II,  $G_1$  is selected from the group consisting of  $-C(O)-R'$ ,  $\alpha$ -amino- $C_{2-11}$  acyl, and  $-C(O)-O-R'$ .  $\alpha$ -Amino- $C_{2-11}$  acyl includes  $\alpha$ -amino acids containing a total of at least 2 carbon atoms and a total of up to 11 carbon atoms, and may also include one or more  
5 heteroatoms selected from the group consisting of O, S, and N.

For certain embodiments, e.g., of Formula III,  $G_2$  is selected from the group consisting of  $-X_2-C(O)-R'$ ,  $\alpha$ -aminoacyl,  $\alpha$ -aminoacyl- $\alpha$ -aminoacyl,  $-X_2-C(O)-O-R'$ ,  $-C(O)-N(R'')R'$ , and  $-S(O)_2-R'$ . For certain of these embodiments,  $X_2$  is selected from the group consisting of a bond;  $-CH_2-O-$ ;  $-CH(CH_3)-O-$ ;  $-C(CH_3)_2-O-$ ; and, in the case of  
10  $-X_2-C(O)-O-R'$ ,  $-CH_2-NH-$ ;

$R'$  and  $R''$  are independently selected from the group consisting of  $C_{1-10}$  alkyl,  $C_{3-7}$  cycloalkyl, phenyl, and benzyl, each of which may be unsubstituted or substituted by one or more substituents independently selected from the group consisting of halogen, hydroxy, nitro, cyano, carboxy,  $C_{1-6}$  alkyl,  $C_{1-4}$  alkoxy, aryl, heteroaryl, aryl- $C_{1-4}$  alkylenyl,  
15 heteroaryl- $C_{1-4}$  alkylenyl, halo- $C_{1-4}$  alkylenyl, halo- $C_{1-4}$  alkoxy,  $-O-C(O)-CH_3$ ,  $-C(O)-O-CH_3$ ,  $-C(O)-NH_2$ ,  $-O-CH_2-C(O)-NH_2$ ,  $-NH_2$ , and  $-S(O)_2-NH_2$ , with the proviso that  $R''$  can also be hydrogen; and

$\alpha$ -aminoacyl is an  $\alpha$ -aminoacyl group derived from an amino acid selected from the group consisting of racemic, D-, and L-amino acids.

20 For certain embodiments, including any one of the above embodiments which include an  $\alpha$ -aminoacyl group,  $\alpha$ -aminoacyl is an  $\alpha$ -aminoacyl group derived from a naturally occurring amino acid selected from the group consisting of racemic, D-, and L-amino acids.

For certain embodiments, including any one of the above embodiments which  
25 include an  $\alpha$ -aminoacyl group,  $\alpha$ -aminoacyl is an  $\alpha$ -aminoacyl group derived from an amino acid found in proteins, wherein the amino acid is selected from the group consisting of racemic, D-, and L-amino acids.

For certain embodiments, including any one of the above embodiments of Formula III,  $G_2$  is selected from the group consisting of  $\alpha$ -amino- $C_{2-5}$  alkanoyl,  $C_{2-6}$  alkanoyl,  
30  $C_{1-6}$  alkoxy carbonyl, and  $C_{1-6}$  alkyl carbamoyl.

For certain embodiments, the hydrogen atom of the 2-hydroxy substituent of Formula II is replaced by  $G_2$ , wherein  $G_2$  is defined as in any one of the above embodiments containing  $G_2$ .

For certain embodiments, including any one of the above embodiments of Formula I, II, or III,  $R_A$  and  $R_B$  are each independently selected from the group consisting of:  
5 hydrogen, halogen, alkenyl, amino,  $-R_{11}$ ,  $-O-R_{11}$ ,  $-S-R_{11}$ , and  $-N(R_{9a})(R_{11})$ .


For certain embodiments, when  $R_A$  and  $R_B$  or either  $R_A$  or  $R_B$  is  $-R_{11}$ ,  $R_{11}$  is selected from the group consisting of alkyl, alkoxyalkylenyl, hydroxyalkylenyl, arylalkylenyl, heteroarylalkylenyl, and heterocyclylalkylenyl, each of which is  
10 unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl; alkoxy; hydroxy; hydroxyalkyl; aryl; aryloxy; arylalkyleneoxy; heteroaryl; heteroaryloxy; heteroarylalkyleneoxy; halogen; haloalkyl; haloalkoxy; mercapto; nitro; cyano; heterocyclyl; amino; alkylamino; dialkylamino; and, in the case of alkyl, heterocyclyl, and heterocyclylalkylenyl, oxo.

For certain embodiments, including any one of the above embodiments of Formula I, II, or III,  $R_A$  and  $R_B$  are independently selected from the group consisting of hydrogen,  $-R_{11}$ ,  $-O-R_{11}$ , and  $-NHR_{11}$ , wherein  $R_{11}$  is alkyl, alkoxyalkylenyl, or hydroxyalkylenyl. For certain of these embodiments,  $R_A$  and  $R_B$  are independently selected from the group consisting of hydrogen,  $C_{1-5}$  alkyl,  $-O-C_{1-4}$  alkyl,  $C_{1-4}$  alkyl- $O-C_{1-4}$  alkylenyl, and  
20  $-NH-C_{1-4}$  alkyl. For certain of these embodiments,  $R_A$  and  $R_B$  are independently selected from the group consisting of hydrogen,  $C_{1-5}$  alkyl,  $-O-C_{1-4}$  alkyl, and  $-NH-C_{1-4}$  alkyl. For certain of these embodiments,  $R_A$  is selected from the group consisting of hydrogen and  $C_{1-5}$  alkyl, and  $R_B$  is selected from the group consisting of  $C_{1-5}$  alkyl,  $-O-C_{1-4}$  alkyl, and  $-NH-C_{1-4}$  alkyl. For certain of these embodiments, except where  $R_A$  and  $R_B$  cannot be  
25 alkyl,  $R_A$  and  $R_B$  are independently hydrogen or alkyl. For certain of these embodiments,  $R_A$  is hydrogen or methyl. For certain of these embodiments,  $R_A$  is hydrogen. For certain of these embodiments,  $R_B$  is  $C_{1-5}$  alkyl. For certain of these embodiments, except where  $R_A$  is hydrogen,  $R_A$  and  $R_B$  are each methyl.

For certain embodiments, including any one of the above embodiments of Formula I, II, or III,  $R_1$  is selected from the group consisting of  $-R_4$ ,  $-X-R_4$ ,  $-X-Y-R_4$ ,  $-X-Y-X-Y-R_4$ ,  
30 and  $-X-R_5$ .

For certain embodiments, including any one of the above embodiments of Formula I, II, or III,  $R_1$  is  $-R_4$  or  $-X-R_4$ .

For certain embodiments, including any one of the above embodiments of Formula I, II, or III,  $R_1$  is selected from the group consisting of aryl- $C_{1-4}$  alkylenyl and heteroaryl- $C_{1-4}$  alkylenyl, wherein the aryl or heteroaryl group is unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl, alkoxy, hydroxyalkyl, haloalkyl, haloalkoxy, halogen, nitro, hydroxy, mercapto, cyano, aryl, aryloxy, arylalkyleneoxy, heteroaryl, heteroaryloxy, heteroarylalkyleneoxy, heterocyclyl, amino, alkylamino, dialkylamino, and (dialkylamino)alkyleneoxy. For certain of these  
 5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55  
 60  
 65  
 70  
 75  
 80  
 85  
 90  
 95  
 100  
 105  
 110  
 115  
 120  
 125  
 130  
 135  
 140  
 145  
 150  
 155  
 160  
 165  
 170  
 175  
 180  
 185  
 190  
 195  
 200  
 205  
 210  
 215  
 220  
 225  
 230  
 235  
 240  
 245  
 250  
 255  
 260  
 265  
 270  
 275  
 280  
 285  
 290  
 295  
 300  
 305  
 310  
 315  
 320  
 325  
 330  
 335  
 340  
 345  
 350  
 355  
 360  
 365  
 370  
 375  
 380  
 385  
 390  
 395  
 400  
 405  
 410  
 415  
 420  
 425  
 430  
 435  
 440  
 445  
 450  
 455  
 460  
 465  
 470  
 475  
 480  
 485  
 490  
 495  
 500  
 505  
 510  
 515  
 520  
 525  
 530  
 535  
 540  
 545  
 550  
 555  
 560  
 565  
 570  
 575  
 580  
 585  
 590  
 595  
 600  
 605  
 610  
 615  
 620  
 625  
 630  
 635  
 640  
 645  
 650  
 655  
 660  
 665  
 670  
 675  
 680  
 685  
 690  
 695  
 700  
 705  
 710  
 715  
 720  
 725  
 730  
 735  
 740  
 745  
 750  
 755  
 760  
 765  
 770  
 775  
 780  
 785  
 790  
 795  
 800  
 805  
 810  
 815  
 820  
 825  
 830  
 835  
 840  
 845  
 850  
 855  
 860  
 865  
 870  
 875  
 880  
 885  
 890  
 895  
 900  
 905  
 910  
 915  
 920  
 925  
 930  
 935  
 940  
 945  
 950  
 955  
 960  
 965  
 970  
 975  
 980  
 985  
 990  
 995  
 1000  
 1005  
 1010  
 1015  
 1020  
 1025  
 1030  
 1035  
 1040  
 1045  
 1050  
 1055  
 1060  
 1065  
 1070  
 1075  
 1080  
 1085  
 1090  
 1095  
 1100  
 1105  
 1110  
 1115  
 1120  
 1125  
 1130  
 1135  
 1140  
 1145  
 1150  
 1155  
 1160  
 1165  
 1170  
 1175  
 1180  
 1185  
 1190  
 1195  
 1200  
 1205  
 1210  
 1215  
 1220  
 1225  
 1230  
 1235  
 1240  
 1245  
 1250  
 1255  
 1260  
 1265  
 1270  
 1275  
 1280  
 1285  
 1290  
 1295  
 1300  
 1305  
 1310  
 1315  
 1320  
 1325  
 1330  
 1335  
 1340  
 1345  
 1350  
 1355  
 1360  
 1365  
 1370  
 1375  
 1380  
 1385  
 1390  
 1395  
 1400  
 1405  
 1410  
 1415  
 1420  
 1425  
 1430  
 1435  
 1440  
 1445  
 1450  
 1455  
 1460  
 1465  
 1470  
 1475  
 1480  
 1485  
 1490  
 1495  
 1500  
 1505  
 1510  
 1515  
 1520  
 1525  
 1530  
 1535  
 1540  
 1545  
 1550  
 1555  
 1560  
 1565  
 1570  
 1575  
 1580  
 1585  
 1590  
 1595  
 1600  
 1605  
 1610  
 1615  
 1620  
 1625  
 1630  
 1635  
 1640  
 1645  
 1650  
 1655  
 1660  
 1665  
 1670  
 1675  
 1680  
 1685  
 1690  
 1695  
 1700  
 1705  
 1710  
 1715  
 1720  
 1725  
 1730  
 1735  
 1740  
 1745  
 1750  
 1755  
 1760  
 1765  
 1770  
 1775  
 1780  
 1785  
 1790  
 1795  
 1800  
 1805  
 1810  
 1815  
 1820  
 1825  
 1830  
 1835  
 1840  
 1845  
 1850  
 1855  
 1860  
 1865  
 1870  
 1875  
 1880  
 1885  
 1890  
 1895  
 1900  
 1905  
 1910  
 1915  
 1920  
 1925  
 1930  
 1935  
 1940  
 1945  
 1950  
 1955  
 1960  
 1965  
 1970  
 1975  
 1980  
 1985  
 1990  
 1995  
 2000  
 2005  
 2010  
 2015  
 2020  
 2025  
 2030  
 2035  
 2040  
 2045  
 2050  
 2055  
 2060  
 2065  
 2070  
 2075  
 2080  
 2085  
 2090  
 2095  
 2100  
 2105  
 2110  
 2115  
 2120  
 2125  
 2130  
 2135  
 2140  
 2145  
 2150  
 2155  
 2160  
 2165  
 2170  
 2175  
 2180  
 2185  
 2190  
 2195  
 2200  
 2205  
 2210  
 2215  
 2220  
 2225  
 2230  
 2235  
 2240  
 2245  
 2250  
 2255  
 2260  
 2265  
 2270  
 2275  
 2280  
 2285  
 2290  
 2295  
 2300  
 2305  
 2310  
 2315  
 2320  
 2325  
 2330  
 2335  
 2340  
 2345  
 2350  
 2355  
 2360  
 2365  
 2370  
 2375  
 2380  
 2385  
 2390  
 2395  
 2400  
 2405  
 2410  
 2415  
 2420  
 2425  
 2430  
 2435  
 2440  
 2445  
 2450  
 2455  
 2460  
 2465  
 2470  
 2475  
 2480  
 2485  
 2490  
 2495  
 2500  
 2505  
 2510  
 2515  
 2520  
 2525  
 2530  
 2535  
 2540  
 2545  
 2550  
 2555  
 2560  
 2565  
 2570  
 2575  
 2580  
 2585  
 2590  
 2595  
 2600  
 2605  
 2610  
 2615  
 2620  
 2625  
 2630  
 2635  
 2640  
 2645  
 2650  
 2655  
 2660  
 2665  
 2670  
 2675  
 2680  
 2685  
 2690  
 2695  
 2700  
 2705  
 2710  
 2715  
 2720  
 2725  
 2730  
 2735  
 2740  
 2745  
 2750  
 2755  
 2760  
 2765  
 2770  
 2775  
 2780  
 2785  
 2790  
 2795  
 2800  
 2805  
 2810  
 2815  
 2820  
 2825  
 2830  
 2835  
 2840  
 2845  
 2850  
 2855  
 2860  
 2865  
 2870  
 2875  
 2880  
 2885  
 2890  
 2895  
 2900  
 2905  
 2910  
 2915  
 2920  
 2925  
 2930  
 2935  
 2940  
 2945  
 2950  
 2955  
 2960  
 2965  
 2970  
 2975  
 2980  
 2985  
 2990  
 2995  
 3000  
 3005  
 3010  
 3015  
 3020  
 3025  
 3030  
 3035  
 3040  
 3045  
 3050  
 3055  
 3060  
 3065  
 3070  
 3075  
 3080  
 3085  
 3090  
 3095  
 3100  
 3105  
 3110  
 3115  
 3120  
 3125  
 3130  
 3135  
 3140  
 3145  
 3150  
 3155  
 3160  
 3165  
 3170  
 3175  
 3180  
 3185  
 3190  
 3195  
 3200  
 3205  
 3210  
 3215  
 3220  
 3225  
 3230  
 3235  
 3240  
 3245  
 3250  
 3255  
 3260  
 3265  
 3270  
 3275  
 3280  
 3285  
 3290  
 3295  
 3300  
 3305  
 3310  
 3315  
 3320  
 3325  
 3330  
 3335  
 3340  
 3345  
 3350  
 3355  
 3360  
 3365  
 3370  
 3375  
 3380  
 3385  
 3390  
 3395  
 3400  
 3405  
 3410  
 3415  
 3420  
 3425  
 3430  
 3435  
 3440  
 3445  
 3450  
 3455  
 3460  
 3465  
 3470  
 3475  
 3480  
 3485  
 3490  
 3495  
 3500  
 3505  
 3510  
 3515  
 3520  
 3525  
 3530  
 3535  
 3540  
 3545  
 3550  
 3555  
 3560  
 3565  
 3570  
 3575  
 3580  
 3585  
 3590  
 3595  
 3600  
 3605  
 3610  
 3615  
 3620  
 3625  
 3630  
 3635  
 3640  
 3645  
 3650  
 3655  
 3660  
 3665  
 3670  
 3675  
 3680  
 3685  
 3690  
 3695  
 3700  
 3705  
 3710  
 3715  
 3720  
 3725  
 3730  
 3735  
 3740  
 3745  
 3750  
 3755  
 3760  
 3765  
 3770  
 3775  
 3780  
 3785  
 3790  
 3795  
 3800  
 3805  
 3810  
 3815  
 3820  
 3825  
 3830  
 3835  
 3840  
 3845  
 3850  
 3855  
 3860  
 3865  
 3870  
 3875  
 3880  
 3885  
 3890  
 3895  
 3900  
 3905  
 3910  
 3915  
 3920  
 3925  
 3930  
 3935  
 3940  
 3945  
 3950  
 3955  
 3960  
 3965  
 3970  
 3975  
 3980  
 3985  
 3990  
 3995  
 4000  
 4005  
 4010  
 4015  
 4020  
 4025  
 4030  
 4035  
 4040  
 4045  
 4050  
 4055  
 4060  
 4065  
 4070  
 4075  
 4080  
 4085  
 4090  
 4095  
 4100  
 4105  
 4110  
 4115  
 4120  
 4125  
 4130  
 4135  
 4140  
 4145  
 4150  
 4155  
 4160  
 4165  
 4170  
 4175  
 4180  
 4185  
 4190  
 4195  
 4200  
 4205  
 4210  
 4215  
 4220  
 4225  
 4230  
 4235  
 4240  
 4245  
 4250  
 4255  
 4260  
 4265  
 4270  
 4275  
 4280  
 4285  
 4290  
 4295  
 4300  
 4305  
 4310  
 4315  
 4320  
 4325  
 4330  
 4335  
 4340  
 4345  
 4350  
 4355  
 4360  
 4365  
 4370  
 4375  
 4380  
 4385  
 4390  
 4395  
 4400  
 4405  
 4410  
 4415  
 4420  
 4425  
 4430  
 4435  
 4440  
 4445  
 4450  
 4455  
 4460  
 4465  
 4470  
 4475  
 4480  
 4485  
 4490  
 4495  
 4500  
 4505  
 4510  
 4515  
 4520  
 4525  
 4530  
 4535  
 4540  
 4545  
 4550  
 4555  
 4560  
 4565  
 4570  
 4575  
 4580  
 4585  
 4590  
 4595  
 4600  
 4605  
 4610  
 4615  
 4620  
 4625  
 4630  
 4635  
 4640  
 4645  
 4650  
 4655  
 4660  
 4665  
 4670  
 4675  
 4680  
 4685  
 4690  
 4695  
 4700  
 4705  
 4710  
 4715  
 4720  
 4725  
 4730  
 4735  
 4740  
 4745  
 4750  
 4755  
 4760  
 4765  
 4770  
 4775  
 4780  
 4785  
 4790  
 4795  
 4800  
 4805  
 4810  
 4815  
 4820  
 4825  
 4830  
 4835  
 4840  
 4845  
 4850  
 4855  
 4860  
 4865  
 4870  
 4875  
 4880  
 4885  
 4890  
 4895  
 4900  
 4905  
 4910  
 4915  
 4920  
 4925  
 4930  
 4935  
 4940  
 4945  
 4950  
 4955  
 4960  
 4965  
 4970  
 4975  
 4980  
 4985  
 4990  
 4995  
 5000  
 5005  
 5010  
 5015  
 5020  
 5025  
 5030  
 5035  
 5040  
 5045  
 5050  
 5055  
 5060  
 5065  
 5070  
 5075  
 5080  
 5085  
 5090  
 5095  
 5100  
 5105  
 5110  
 5115  
 5120  
 5125  
 5130  
 5135  
 5140  
 5145  
 5150  
 5155  
 5160  
 5165  
 5170  
 5175  
 5180  
 5185  
 5190  
 5195  
 5200  
 5205  
 5210  
 5215  
 5220  
 5225  
 5230  
 5235  
 5240  
 5245  
 5250  
 5255  
 5260  
 5265  
 5270  
 5275  
 5280  
 5285  
 5290  
 5295  
 5300  
 5305  
 5310  
 5315  
 5320  
 5325  
 5330  
 5335  
 5340  
 5345  
 5350  
 5355  
 5360  
 5365  
 5370  
 5375  
 5380  
 5385  
 5390  
 5395  
 5400  
 5405  
 5410  
 5415  
 5420  
 5425  
 5430  
 5435  
 5440  
 5445  
 5450  
 5455  
 5460  
 5465  
 5470  
 5475  
 5480  
 5485  
 5490  
 5495  
 5500  
 5505  
 5510  
 5515  
 5520  
 5525  
 5530  
 5535  
 5540  
 5545  
 5550  
 5555  
 5560  
 5565  
 5570  
 5575  
 5580  
 5585  
 5590  
 5595  
 5600  
 5605  
 5610  
 5615  
 5620  
 5625  
 5630  
 5635  
 5640  
 5645  
 5650  
 5655  
 5660  
 5665  
 5670  
 5675  
 5680  
 5685  
 5690  
 5695  
 5700  
 5705  
 5710  
 5715  
 5720  
 5725  
 5730  
 5735  
 5740  
 5745  
 5750  
 5755  
 5760  
 5765  
 5770  
 5775  
 5780  
 5785  
 5790  
 5795  
 5800  
 5805  
 5810  
 5815  
 5820  
 5825  
 5830  
 5835  
 5840  
 5845  
 5850  
 5855  
 5860  
 5865  
 5870  
 5875  
 5880  
 5885  
 5890  
 5895  
 5900  
 5905  
 5910  
 5915  
 5920  
 5925  
 5930  
 5935  
 5940  
 5945  
 5950  
 5955  
 5960  
 5965  
 5970  
 5975  
 5980  
 5985  
 5990  
 5995  
 6000  
 6005  
 6010  
 6015  
 6020  
 6025  
 6030  
 6035  
 6040  
 6045  
 6050  
 6055  
 6060  
 6065  
 6070  
 6075  
 6080  
 6085  
 6090  
 6095  
 6100  
 6105  
 6110  
 6115  
 6120  
 6125  
 6130  
 6135  
 6140  
 6145  
 6150  
 6155  
 6160  
 6165  
 6170  
 6175  
 6180  
 6185  
 6190  
 6195  
 6200  
 6205  
 6210  
 6215  
 6220  
 6225  
 6230  
 6235  
 6240  
 6245  
 6250  
 6255  
 6260  
 6265  
 6270  
 6275  
 6280  
 6285  
 6290  
 6295  
 6300  
 6305  
 6310  
 6315  
 6320  
 6325  
 6330  
 6335  
 6340  
 6345  
 6350  
 6355  
 6360  
 6365  
 6370  
 6375  
 6380  
 6385  
 6390  
 6395  
 6400  
 6405  
 6410  
 6415  
 6420  
 6425  
 6430  
 6435  
 6440  
 6445  
 6450  
 6455  
 6460  
 6465  
 6470  
 6475  
 6480  
 6485  
 6490  
 6495  
 6500  
 6505  
 6510  
 6515  
 6520  
 6525  
 6530  
 6535  
 6540  
 6545  
 6550  
 6555  
 6560  
 6565  
 6570  
 6575  
 6580  
 6585  
 6590  
 6595  
 6600  
 6605  
 6610  
 6615  
 6620  
 6625  
 6630  
 6635  
 6640  
 6645  
 6650  
 6655  
 6660  
 6665  
 6670  
 6675  
 6680  
 6685  
 6690  
 6695  
 6700  
 6705  
 6710  
 6715  
 6720  
 6725  
 6730  
 6735  
 6740  
 6745  
 6750  
 6755  
 6760  
 6765  
 6770  
 6775

embodiments, R<sub>1</sub> is . Alternatively, for certain of these  
embodiments, R<sub>1</sub> is -C<sub>2-5</sub> alkylenyl-NH-Q-R<sub>4</sub>. For certain of these embodiments where Q  
is present, Q is -C(O)-, S(O)<sub>2</sub>-, or -C(O)-NH-, and R<sub>4</sub> is C<sub>1-6</sub>alkyl.

For certain embodiments, including any one of the above embodiments of Formula  
5 I, II, or III, where not excluded, R<sub>1</sub> is selected from the group consisting of -N(R<sub>1</sub>')-Q-R<sub>4</sub>,  
-N(R<sub>1</sub>')-X<sub>1</sub>-Y<sub>1</sub>-R<sub>4</sub>, and -N(R<sub>1</sub>')-X<sub>1</sub>-R<sub>5a</sub>.

For certain embodiments, including any one of the above embodiments of Formula  
I, II, or III, where not excluded, R<sub>1</sub> is -N(R<sub>1</sub>')-Q-R<sub>4</sub>. For certain of these embodiments, R<sub>1</sub>'  
is hydrogen, Q is a bond, and R<sub>4</sub> is aryl, heteroaryl, aryl-C<sub>1-3</sub> alkylenyl, or  
10 heteroaryl-C<sub>1-3</sub> alkylenyl.

For certain embodiments, R<sub>1</sub>' is selected from the group consisting of hydrogen,  
C<sub>1-20</sub> alkyl, hydroxy-C<sub>2-20</sub> alkylenyl, and alkoxy-C<sub>2-20</sub> alkylenyl.

For certain embodiments, R<sub>1</sub>' is hydrogen or methyl.

For certain embodiments, R<sub>1</sub>' is hydrogen.

15 For certain embodiments, R<sub>4</sub> is selected from the group consisting of hydrogen,  
alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl,  
heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl  
wherein the alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl,  
heteroaryl, heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and  
20 heterocyclyl groups can be unsubstituted or substituted by one or more substituents  
independently selected from the group consisting of alkyl; alkoxy; hydroxyalkyl;  
haloalkyl; haloalkoxy; halogen; nitro; hydroxy; mercapto; cyano; aryl; aryloxy;  
arylalkyleneoxy; heteroaryl; heteroaryloxy; heteroarylalkyleneoxy; heterocyclyl; amino;  
alkylamino; dialkylamino; (dialkylamino)alkyleneoxy; and, in the case of alkyl, alkenyl,  
25 alkynyl, and heterocyclyl, oxo.

For certain embodiments, R<sub>4</sub> is selected from the group consisting of  
aryl-C<sub>1-4</sub> alkylenyl and heteroaryl-C<sub>1-4</sub> alkylenyl, wherein the aryl or heteroaryl group is  
unsubstituted or substituted by one or more substituents independently selected from the  
group consisting of alkyl, alkoxy, hydroxyalkyl, haloalkyl, haloalkoxy, halogen, nitro,  
30 hydroxy, mercapto, cyano, aryl, aryloxy, arylalkyleneoxy, heteroaryl, heteroaryloxy,

heteroarylalkyleneoxy, heterocyclyl, amino, alkylamino, dialkylamino, and (dialkylamino)alkyleneoxy.

For certain embodiments,  $R_4$  is benzyl, which is unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl, alkoxy, haloalkyl, haloalkoxy, and halogen.

For certain embodiments,  $R_4$  is benzyl.

For certain embodiments,  $R_4$  is tetrahydro-2*H*-pyran-4-ylmethyl.

For certain embodiments,  $R_4$  is aryl, heteroaryl, aryl- $C_{1-3}$  alkylenyl, or heteroaryl- $C_{1-3}$  alkylenyl.


For certain embodiments,  $R_4$  is isoxazol-3-yl, isoxazol-5-yl, or thiazol-2-yl, each of which is unsubstituted or substituted by methyl or 4-fluorophenyl.

For certain embodiments,  $R_4$  is phenyl.


For certain embodiments,  $R_4$  is  $C_{1-6}$  alkyl.

For certain embodiments,  $R_4$  is  $C_{1-3}$  alkyl.

For certain embodiments,  $R_{5a}$  is selected from the group consisting of:




For certain embodiments,  $R_{5a}$  is




For certain embodiments,  $R_{5a}$  is

For certain embodiments,  $R_5$  is selected from the group consisting of:



For certain embodiments,  $R_5$  is



For certain embodiments,  $R_6$  is selected from the group consisting of =O and =S.

For certain embodiments,  $R_6$  is =O.

For certain embodiments,  $R_7$  is  $C_{2-7}$  alkylene.

For certain embodiments,  $R_7$  is  $C_{2-4}$  alkylene.

5 For certain embodiments,  $R_7$  is ethylene.

For certain embodiments,  $R_8$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl, hydroxy- $C_{1-10}$  alkylenyl,  $C_{1-10}$  alkoxy- $C_{1-10}$  alkylenyl, aryl- $C_{1-10}$  alkylenyl, and heteroaryl- $C_{1-10}$  alkylenyl.

For certain embodiments,  $R_8$  is hydrogen or  $C_{1-4}$  alkyl.

10 For certain embodiments,  $R_8$  is hydrogen.

For certain embodiments,  $R_{9a}$  is selected from the group consisting of hydrogen and  $C_{1-4}$  alkyl.

For certain embodiments,  $R_{9a}$  is hydrogen.

15 For certain embodiments,  $R_9$  is selected from the group consisting of hydrogen and alkyl.

For certain embodiments,  $R_{10}$  is  $C_{3-8}$  alkylene.

For certain embodiments,  $R_{10}$  is pentylene.

20 For certain embodiments,  $R_{11}$  is selected from the group consisting of alkyl, alkoxyalkylenyl, hydroxyalkylenyl, aryl, arylalkylenyl, heteroaryl, heteroarylalkylenyl, heterocyclyl, and heterocyclylalkylenyl, each of which is unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl; alkoxy; hydroxy; hydroxyalkyl; aryl; aryloxy; arylalkyleneoxy; heteroaryl; heteroaryloxy; heteroarylalkyleneoxy; halogen; haloalkyl; haloalkoxy; mercapto; nitro; cyano; heterocyclyl; amino; alkylamino; dialkylamino; and, in the case of alkyl, heterocyclyl, and  
25 heterocyclylalkylenyl, oxo.

For certain embodiments,  $R_{11}$  is selected from the group consisting of alkyl, alkoxyalkylenyl, hydroxyalkylenyl, arylalkylenyl, heteroarylalkylenyl, and heterocyclylalkylenyl, each of which is unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl; alkoxy; hydroxy; hydroxyalkyl; aryl; aryloxy; arylalkyleneoxy; heteroaryl; heteroaryloxy; heteroarylalkyleneoxy; halogen; haloalkyl; haloalkoxy; mercapto; nitro; cyano;  
30 heteroarylalkyleneoxy; halogen; haloalkyl; haloalkoxy; mercapto; nitro; cyano;

heterocyclyl; amino; alkylamino; dialkylamino; and, in the case of alkyl, and heterocyclylalkylenyl, oxo.

For certain embodiments,  $R_{11}$  is alkyl, alkoxyalkylenyl, or hydroxyalkylenyl.

For certain embodiments,  $R_{11}$  is pentyl.

5 For certain embodiments, A is selected from the group consisting of  $-CH_2-$ ,  $-O-$ ,  $-C(O)-$ ,  $-S(O)_{0-2}-$ , and  $-N(-Q-R_4)-$ . For certain embodiments, A is  $-O-$ .

For certain embodiments, A' is selected from the group consisting of  $-O-$ ,  $-S(O)_{0-2}-$ ,  $-N(-Q-R_4)-$ , and  $-CH_2-$ .

10 For certain embodiments, Q is selected from the group consisting of a bond,  $-C(R_6)-$ ,  $-C(R_6)-C(R_6)-$ ,  $-S(O)_2-$ ,  $-C(R_6)-N(R_8)-W-$ ,  $-S(O)_2-N(R_8)-$ ,  $-C(R_6)-O-$ ,  $-C(R_6)-S-$ , and  $-C(R_6)-N(OR_9)-$ . For certain embodiments, Q is  $-C(R_6)-N(R_8)-$ ,  $-C(R_6)-$ , or  $-S(O)_2-$ . For certain embodiments, Q is  $-C(O)-N(H)-$ ,  $-C(O)-$ , or  $-S(O)_2-$ . For certain embodiments, Q is  $-C(R_6)-N(R_8)-$ . For certain embodiments, Q is  $-C(O)-NH-$ . For certain embodiments, Q is  $-S(O)_2-$ . For certain embodiments, Q is  $-C(R_6)-$ . For certain embodiments, Q is  $-C(O)-$ . For certain embodiments, Q is a bond.


For certain embodiments, V is selected from the group consisting of  $-C(R_6)-$ ,  $-O-C(R_6)-$ ,  $-N(R_8)-C(R_6)-$ , and  $-S(O)_2-$ . For certain embodiments, V is  $-N(R_8)-C(O)-$ .

For certain embodiments, W is selected from the group consisting of a bond,  $-C(O)-$ , and  $-S(O)_2-$ . For certain embodiments, W is a bond.

20 For certain embodiments, X is selected from the group consisting of alkylene, alkenylene, alkynylene, arylene, heteroarylene, and heterocyclylene, wherein the alkylene, alkenylene, and alkynylene groups can be optionally interrupted or terminated by arylene, heteroarylene or heterocyclylene and optionally interrupted by one or more  $-O-$  groups.

For certain embodiments, X is  $C_{1-4}$  alkylene.

25 For certain embodiments,  $-X-$  is



For certain embodiments, X is methylene.



For certain embodiments, X is  $-C_{1-4}$  alkylene- $O-C_{1-4}$  alkylene-.

For certain of these embodiments, X is  $-(CH_2)_2-O-(CH_2)_3-$ .

30 For certain embodiments,  $X_1$  is  $C_{2-20}$  alkylene. For certain embodiments,  $X_1$  is  $C_{2-4}$  alkylene.




For certain embodiments, Y is selected from the group consisting of -O-,  
 -S(O)<sub>0.2</sub>-, -S(O)<sub>2</sub>-N(R<sub>8</sub>)-, -C(R<sub>6</sub>)-, -C(R<sub>6</sub>)-O-, -O-C(R<sub>6</sub>)-, -O-C(O)-O-, -N(R<sub>8</sub>)-Q-,  
 -C(R<sub>6</sub>)-N(R<sub>8</sub>)-, -O-C(R<sub>6</sub>)-N(R<sub>8</sub>)-, -C(R<sub>6</sub>)-N(OR<sub>9</sub>)-, -O-N(R<sub>8</sub>)-Q-, -O-N=C(R<sub>4</sub>)-,



5

For certain embodiments, Y is -N(R<sub>8</sub>)-C(O)-, -N(R<sub>8</sub>)-S(O)<sub>2</sub>-,  
 -N(R<sub>8</sub>)-C(R<sub>6</sub>)-N(R<sub>8</sub>)-, -N(R<sub>8</sub>)-C(R<sub>6</sub>)-N(R<sub>8</sub>)-C(O)-, -N(R<sub>8</sub>)-C(R<sub>6</sub>)-O-,




For certain embodiments, Y is -S(O)<sub>2</sub>-.


10

For certain embodiments, Y is -NH-Q-.


For certain embodiments, Y is



For certain embodiments, Y is



For certain embodiments, Y<sub>1</sub> is selected from the group consisting of -O-,  
 -S(O)<sub>0.2</sub>-, -S(O)<sub>2</sub>-N(R<sub>8</sub>)-, -N(R<sub>8</sub>)-Q-, -C(R<sub>6</sub>)-N(R<sub>8</sub>)-, -O-C(R<sub>6</sub>)-N(R<sub>8</sub>)-, and



15

For certain embodiments, a and b are independently integers from 1 to 6 with the  
 proviso that a + b is ≤ 7. For certain embodiments, a and b are each 2.

For certain embodiments, the present invention provides a pharmaceutical  
 composition comprising a therapeutically effective amount of a compound or salt of any  
 one of the above embodiments of Formulas I, II, and III, and a pharmaceutically  
 acceptable carrier.

20

For certain embodiments, the present invention provides a method of inducing cytokine biosynthesis in an animal comprising administering an effective amount of a compound or salt of any one of the above embodiments of Formulas I, II, and III, or a pharmaceutical composition comprising an effective amount of any one of the above  
5       embodiments of Formulas I, II, and III to the animal. For certain of these embodiments, the cytokine is selected from the group consisting of IFN- $\alpha$ , TNF- $\alpha$ , IL-6, and IL-10. For certain of these embodiments, the cytokine is IFN- $\alpha$  or IFN- $\alpha$  and TNF- $\alpha$ . For certain of these embodiments, the cytokine is IFN- $\alpha$ .

For certain embodiments, the present invention provides a method of selectively  
10       inducing the biosynthesis of IFN- $\alpha$  in an animal comprising administering an effective amount of a compound or salt of any one of the above embodiments of Formulas I, II, and III, or a pharmaceutical composition comprising an effective amount of any one of the above embodiments of Formulas I, II, and III to the animal.

For certain embodiments, the present invention provides a method of treating a  
15       viral disease in an animal comprising administering a therapeutically effective amount of a compound or salt of any one of the above embodiments of Formulas I, II, and III, or a pharmaceutical composition comprising a therapeutically effective amount of any one of the above embodiments of Formulas I, II, and III to the animal.

For certain embodiments, the present invention provides a method of treating a  
20       viral disease in an animal comprising administering a therapeutically effective amount of a compound or salt of any one of the above embodiments of Formulas I, II, and III, or a pharmaceutical composition comprising a therapeutically effective amount of any one of the above embodiments of Formulas I, II, and III to the animal; and selectively inducing the biosynthesis of IFN- $\alpha$  in the animal.

For certain embodiments, the present invention provides a method of treating a  
25       neoplastic disease in an animal comprising administering a therapeutically effective amount of a compound or salt of any one of the above embodiments of Formulas I, II, and III, or a pharmaceutical composition comprising a therapeutically effective amount of any one of the above embodiments of Formulas I, II, and III to the animal.

For certain embodiments, the present invention provides a method of treating a  
30       neoplastic disease in an animal comprising administering a therapeutically effective amount of a compound or salt of any one of the above embodiments of Formulas I, II, and

III, or a pharmaceutical composition comprising a therapeutically effective amount of any one of the above embodiments of Formulas I, II, and III to the animal; and selectively inducing the biosynthesis of IFN- $\alpha$  in the animal.

As used herein, the terms "alkyl", "alkenyl", "alkynyl" and the prefix "alk-" are inclusive of both straight chain and branched chain groups and of cyclic groups, e.g., cycloalkyl and cycloalkenyl. Unless otherwise specified, these groups contain from 1 to 20 carbon atoms, with alkenyl groups containing from 2 to 20 carbon atoms, and alkynyl groups containing from 2 to 20 carbon atoms. In some embodiments, these groups have a total of up to 10 carbon atoms, up to 8 carbon atoms, up to 6 carbon atoms, or up to 4 carbon atoms. Cyclic groups can be monocyclic or polycyclic and preferably have from 3 to 10 ring carbon atoms. Exemplary cyclic groups include cyclopropyl, cyclopropylmethyl, cyclobutyl, cyclobutylmethyl, cyclopentyl, cyclopentylmethyl, cyclohexyl, cyclohexylmethyl, adamantyl, and substituted and unsubstituted bornyl, norbornyl, and norbornenyl.

Unless otherwise specified, "alkylene", "-alkylene-", "alkenylene", "-alkenylene-", "alkynylene", and "-alkynylene-" are the divalent forms of the "alkyl", "alkenyl", and "alkynyl" groups defined above. The terms "alkylenyl", "alkenylenyl", and "alkynylenyl" are used when "alkylene", "alkenylene", and "alkynylene", respectively, are substituted. For example, an arylalkylenyl group comprises an "alkylene" moiety to which an aryl group is attached.

The term "haloalkyl" is inclusive of alkyl groups that are substituted by one or more halogen atoms, including perfluorinated groups. This is also true of other groups that include the prefix "halo-". Examples of suitable haloalkyl groups are chloromethyl, trifluoromethyl, and the like.

The term "aryl" as used herein includes carbocyclic aromatic rings or ring systems. Examples of aryl groups include phenyl, naphthyl, biphenyl, fluorenyl and indenyl.

Unless otherwise indicated, the term "heteroatom" refers to the atoms O, S, or N.

The term "heteroaryl" includes aromatic rings or ring systems that contain at least one ring heteroatom (e.g., O, S, N). In some embodiments, the term "heteroaryl" includes a ring or ring system that contains 2 to 12 carbon atoms, 1 to 3 rings, 1 to 4 heteroatoms, and O, S, and/or N as the heteroatoms. Suitable heteroaryl groups include furyl, thienyl, pyridyl, quinolinyl, isoquinolinyl, indolyl, isoindolyl, triazolyl, pyrrolyl, tetrazolyl,

imidazolyl, pyrazolyl, oxazolyl, thiazolyl, benzofuranyl, benzothiophenyl, carbazolyl, benzoxazolyl, pyrimidinyl, benzimidazolyl, quinoxaliny, benzothiazolyl, naphthyridinyl, isoxazolyl, isothiazolyl, purinyl, quinazolinyl, pyrazinyl, 1-oxidopyridyl, pyridazinyl, triazinyl, tetrazinyl, oxadiazolyl, thiadiazolyl, and so on.

The term "heterocyclyl" includes non-aromatic rings or ring systems that contain at least one ring heteroatom (e.g., O, S, N) and includes all of the fully saturated and partially unsaturated derivatives of the above mentioned heteroaryl groups. In some embodiments, the term "heterocyclyl" includes a ring or ring system that contains 2 to 12 carbon atoms, 1 to 3 rings, 1 to 4 heteroatoms, and O, S, and N as the heteroatoms. Exemplary heterocyclyl groups include pyrrolidinyl, tetrahydrofuranyl, morpholinyl, thiomorpholinyl, 1,1-dioxothiomorpholinyl, piperidinyl, piperazinyl, thiazolidinyl, imidazolidinyl, isothiazolidinyl, tetrahydropyranyl, quinuclidinyl, homopiperidinyl (azepanyl), 1,4-oxazepanyl, homopiperazinyl (diazepanyl), 1,3-dioxolanyl, aziridinyl, azetidiny, dihydroisoquinolin-(1*H*)-yl, octahydroisoquinolin-(1*H*)-yl, dihydroquinolin-(2*H*)-yl, octahydroquinolin-(2*H*)-yl, dihydro-1*H*-imidazolyl, 3-azabicyclo[3.2.2]non-3-yl, and the like.


The term "heterocyclyl" includes bicyclic and tricyclic heterocyclic ring systems. Such ring systems include fused and/or bridged rings and spiro rings. Fused rings can include, in addition to a saturated or partially saturated ring, an aromatic ring, for example, a benzene ring. Spiro rings include two rings joined by one spiro atom and three rings joined by two spiro atoms.

When "heterocyclyl" contains a nitrogen atom, the point of attachment of the heterocyclyl group may be the nitrogen atom.

25 The terms "arylene", "heteroarylene", and "heterocyclylene" are the divalent forms of the "aryl", "heteroaryl", and "heterocyclyl" groups defined above. The terms, "arylenyl", "heteroarylenyl", and "heterocyclylenyl" are used when "arylene", "heteroarylene", and "heterocyclylene", respectively, are substituted. For example, an alkylarylenyl group comprises an arylene moiety to which an alkyl group is attached.

When a group (or substituent or variable) is present more than once in any Formula  
30 described herein, each group (or substituent or variable) is independently selected, whether

explicitly stated or not. For example, for the formula



each  $R_7$  group

is independently selected. In another example, when more than one Y group is present, each Y group is independently selected. In a further example, when more than one -N(R<sub>8</sub>)-C(R<sub>6</sub>)-N(R<sub>8</sub>)- group is present (e.g., more than one Y group is present, and both contain a -N(R<sub>8</sub>)-C(R<sub>6</sub>)-N(R<sub>8</sub>)- group) each R<sub>8</sub> group is independently selected and each R<sub>6</sub> group is independently selected.

The invention is inclusive of the compounds described herein (including intermediates) in any of their pharmaceutically acceptable forms, including isomers (e.g., diastereomers and enantiomers), salts, solvates, polymorphs, prodrugs, and the like. In particular, if a compound is optically active, the invention specifically includes each of the compound's enantiomers as well as racemic and scalemic mixtures of the enantiomers. It should be understood that the term "compound" includes any or all of such forms, whether explicitly stated or not (although at times, "salts" are explicitly stated).

The term "prodrug" means a compound that can be transformed in vivo to yield an immune response modifying compound, including any of the salt, solvated, polymorphic, or isomeric forms described above. The prodrug, itself, may be an immune response modifying compound, including any of the salt, solvated, polymorphic, or isomeric forms described above. The transformation may occur by various mechanisms, such as through a chemical (e.g., solvolysis or hydrolysis, for example, in the blood) or enzymatic biotransformation. A discussion of the use of prodrugs is provided by T. Higuchi and W. Stella, "Pro-drugs as Novel Delivery Systems," Vol. 14 of the A. C. S. Symposium Series, and in *Bioreversible Carriers in Drug Design*, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.

#### Preparation of the Compounds

Compounds of the invention may be synthesized by synthetic routes that include processes analogous to those well known in the chemical arts, particularly in light of the description contained herein. The starting materials are generally available from commercial sources such as Aldrich Chemicals (Milwaukee, Wisconsin, USA) or are readily prepared using methods well known to those skilled in the art (e.g. prepared by methods generally described in Louis F. Fieser and Mary Fieser, *Reagents for Organic Synthesis*, v. 1-19, Wiley, New York, (1967-1999 ed.); Alan R. Katritzky, Otto Meth-Cohn, Charles W. Rees, *Comprehensive Organic Functional Group Transformations*, v 1-

6, Pergamon Press, Oxford, England, (1995); Barry M. Trost and Ian Fleming, *Comprehensive Organic Synthesis*, v. 1-8, Pergamon Press, Oxford, England, (1991); or *Beilsteins Handbuch der organischen Chemie*, 4, Aufl. Ed. Springer-Verlag, Berlin, Germany, including supplements (also available via the Beilstein online database)).

5 For illustrative purposes, the reaction schemes depicted below provide potential routes for synthesizing the compounds of the present invention as well as key intermediates. For more detailed description of the individual reaction steps, see the EXAMPLES section below. Those skilled in the art will appreciate that other synthetic routes may be used to synthesize the compounds of the invention. Although specific  
10 starting materials and reagents are depicted in the reaction schemes and discussed below, other starting materials and reagents can be easily substituted to provide a variety of derivatives and/or reaction conditions. In addition, many of the compounds prepared by the methods described below can be further modified in light of this disclosure using conventional methods well known to those skilled in the art.

15 In the preparation of compounds of the invention it may sometimes be necessary to protect a particular functionality while reacting other functional groups on an intermediate. The need for such protection will vary depending on the nature of the particular functional group and the conditions of the reaction step. Suitable amino protecting groups include acetyl, trifluoroacetyl, *tert*-butoxycarbonyl (Boc), benzyloxycarbonyl, and 9-  
20 fluorenylmethoxycarbonyl (Fmoc). Suitable hydroxy protecting groups include acetyl and silyl groups such as the *tert*-butyl dimethylsilyl group. For a general description of protecting groups and their use, see T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons, New York, USA, 1991.

25 Conventional methods and techniques of separation and purification can be used to isolate compounds of the invention or pharmaceutically acceptable salts thereof, as well as various intermediates related thereto. Such techniques may include, for example, all types of chromatography (high performance liquid chromatography (HPLC), column chromatography using common absorbents such as silica gel, and thin layer chromatography, recrystallization, and differential (i.e., liquid-liquid) extraction  
30 techniques.

Compounds of the invention can be prepared according to Reaction Scheme I, wherein R<sub>A</sub>, R<sub>B</sub>, G<sub>1</sub>, G<sub>2</sub>, and R<sub>1</sub> are as defined above, and Bn is benzyl, *p*-methoxybenzyl,

*p*-methylbenzyl, or 2-furanylmethyl. In step (1) of Reaction Scheme I, a 2,4-dichloro-3-nitropyridine of Formula V is reacted with an amine of Formula  $R_1-NH_2$ . The reaction can be conveniently carried out by adding the amine to a solution of a compound of Formula V in the presence of a base such as triethylamine. The reaction is carried out in a suitable solvent, such as dichloromethane, chloroform, or *N,N*-dimethylformamide (DMF) and may be carried out at room temperature, a sub-ambient temperature such as 0 ° C, or an elevated temperature such as the reflux temperature of the solvent. Many 2,4-dichloro-3-nitropyridines of Formula V are known or can be prepared by known methods; see, for example, U.S. Patent No. 6,525,064 (Dellaria *et al.*). For example, they are readily prepared by chlorinating 4-hydroxy-3-nitro-2(1*H*)-pyridones with a chlorinating agent such as phosphorus(III) oxychloride. Many 4-hydroxy-3-nitro-2(1*H*)-pyridones are known or can be prepared by known methods; see, for example, U.S. Patent No. 5,446,153 (Lindstrom *et al.*) and the references cited therein. Other 2,4-dichloro-3-nitropyridines of Formula V can be prepared according to methods described in Reaction Scheme II.

Numerous amines of Formula  $R_1-NH_2$  are commercially available; others can be prepared by known methods. For example, a variety of substituted and unsubstituted alkyl and arylalkylenyl amines, isomeric (aminomethyl)pyridines, and alkyl, aryl, or arylalkylenyl hydrazines or hydrazine salts are commercially available. In certain preferred embodiments,  $R_1$  is a (5-substituted-isoxazol-3-yl)methyl group. (5-Substituted-isoxazol-3-yl)methylamines can be prepared by the following four-step method. In part (i), a protected amino-aldehyde of formula  $(PG)_2-X-CH=O$ , wherein PG is a nitrogen protecting group and X is as defined above, is converted to an aldoxime of formula  $(PG)_2-X-CH=N-OH$  using conventional methods. For example, an aldehyde can be combined with hydroxylamine hydrochloride in the presence of base such as triethylamine in a suitable solvent such as dichloromethane. The reaction can be run at room temperature. Protected amino-aldehydes can be prepared using conventional methods. For example, phthalimidoacetaldehyde diethyl acetal is a commercially available compound that can be treated with acid to provide an aldehyde of formula  $(PG)_2-X-CH=O$ . In part (ii), an aldoxime of formula  $(PG)_2-X-CH=N-OH$  is converted to an  $\alpha$ -chloraldoxime of formula  $(PG)_2-X-C(Cl)=N-OH$  by treatment with *N*-chlorosuccinimide in a suitable solvent such as DMF. The reaction may be carried out

initially below room temperature, at 0 °C for example, and then heated at an elevated temperature in the range of 40 °C to 50 °C. The  $\alpha$ -chloroaldoxime of formula (PG)<sub>2</sub>-X-C(Cl)=N-OH can optionally be isolated before it is converted in part (iii) to a protected (5-substituted-isoxazol-3-yl)methylamine by treatment with a base such as triethylamine to generate a nitrile oxide in the presence of an alkyne in a suitable solvent such as dichloromethane at room temperature. The nitrile oxide and alkyne undergo a [3+2] cycloaddition reaction to provide a protected (5-substituted-isoxazol-3-yl)methylamine, which is then deprotected in part (iv) using conventional methods. When a phthalimide protecting group is used, the deprotection can be carried out by combining the phthalimide-protected (5-substituted-isoxazol-3-yl)methylamine with hydrazine or hydrazine hydrate in a suitable solvent such as ethanol or solvent mixture such as ethanol/THF. The deprotection reaction can be carried out at room temperature or at an elevated temperature such as the reflux temperature of the solvent.

Some amines of the Formula H<sub>2</sub>N-R<sub>1</sub> can be made according to the following methods. For some embodiments, R<sub>1</sub> is a 1-hydroxycycloalkylmethyl group, a (4-hydroxytetrahydro-2H-pyran-4-yl)methyl group, or a group derived from a [1-(*tert*-butoxycarbonyl)-4-hydroxypiperidin-4-yl]methyl group. The corresponding amines of Formula H<sub>2</sub>N-R<sub>1</sub> can be prepared by combining a cyclic ketone, such as cyclopentanone, cyclobutanone, tetrahydro-4H-pyran-4-one, and *tert*-butyl 4-oxo-1-piperidinecarboxylate, with excess nitromethane in a suitable solvent such as ethanol or methanol in the presence of a catalytic amount of base such as sodium ethoxide or sodium hydroxide and reducing the resultant nitromethyl-substituted compound using conventional heterogeneous hydrogenation conditions. The hydrogenation is typically carried out in the presence of a catalyst such as palladium hydroxide on carbon, palladium on carbon, or Raney nickel in a suitable solvent such as ethanol. Both the reaction with nitromethane and the reduction can be carried out at room temperature. A wide variety of cyclic ketones can be obtained from commercial sources; others can be synthesized using known synthetic methods.

In step (2) of Reaction Scheme I, the chloro group in a pyridine of Formula VI is displaced by an amine of Formula HN(Bn)<sub>2</sub> to provide a pyridine of Formula VII. The displacement is conveniently carried out by combining an amine of Formula HN(Bn)<sub>2</sub> and a compound of Formula VI in a suitable solvent such as toluene or xylenes in the presence




of a base such as triethylamine and heating at an elevated temperature such as the reflux temperature of the solvent.

In step (3) of Reaction Scheme I, a compound of Formula VII is reduced to provide a pyridine-2,3,4-triamine of Formula VIII. The reduction can be carried out using nickel boride, prepared *in situ* from sodium borohydride and nickel(II) chloride. The reduction is conveniently carried out by adding a solution of a pyridine of Formula VII in a suitable solvent or solvent mixture such as dichloromethane/methanol to a mixture of excess sodium borohydride and catalytic or stoichiometric nickel(II) chloride in methanol. The reaction can be carried out at room temperature.

In step (4) of Reaction Scheme I, a pyridine-2,3,4-triamine of Formula VIII is cyclized to provide a 1*H*-imidazo[4,5-*c*]pyridin-2-ol of Formula IX. The cyclization can be conveniently carried out by heating a pyridine-2,3,4-triamine of Formula VIII with carbonyl diimidazole in a suitable solvent such as tetrahydrofuran (THF), *tert*-butyl methyl ether, dichloromethane, or DMF. The reaction may be carried out at room temperature or, preferably, at an elevated temperature such as the reflux temperature of the solvent.


In step (5) of Reaction Scheme I, the protecting groups are removed from the 4-amine of a 1*H*-imidazo[4,5-*c*]pyridin-2-ol of Formula IX to provide a 1*H*-imidazo[4,5-*c*]pyridin-2-ol of Formula I. For certain embodiments, the deprotection can be conveniently carried out on a Parr apparatus under hydrogenolysis conditions using a suitable heterogeneous catalyst such as palladium on carbon in a solvent such as ethanol. Alternatively, when Bn is *p*-methoxybenzyl, step (5) may be carried out by combining trifluoroacetic acid and a compound of Formula IX and stirring at room temperature or heating at an elevated temperature such as 50 °C to 70 °C.

Certain amines of Formula  $R_1-NH_2$  provide a compound of Formula VI that contain a functional group or protected functional group that can be transformed in a subsequent step to provide compounds of Formula I with a variety of different  $R_1$  groups. For example, protected diamines of Formula Boc-N( $R_8$ )-X-NH<sub>2</sub>,




are commercially available or can be prepared by known methods; see, for example, U.S. Patent No. 6,797,718 (Dellaria *et al.*) and Carceller, E. *et al.*, *J. Med. Chem.*, 39, pp.487-493 (1996). The Boc-protected amino

group can be subjected to the reaction conditions of steps (2) through (4) of Reaction Scheme I. The Boc-protecting group may be removed in step (5) if the acidic conditions are used, or it can be removed by conventional methods after step (5). The resulting compound of Formula I having an  $-X-N(R_8)H$ ,



group at the  $R_1$  position can be converted to a compound having an  $-X-N(R_8)-Q-R_4$ ,



group using known methods; see, for example, Dellaria et al. in U.S. Patent Nos. 6,525,064, 6,545,016, 6,545,017, and 6,797,718. In other examples, amino esters of Formula  $H_2N-X-C(O)-O\text{-alkyl}$  or hydrochloride salts thereof can be used in step (1), and the resulting compound of Formula VI can be converted in subsequent steps to a compound of Formula I having an  $-X-C(R_6)-R_4$ ,  $-X-C(R_6)-N(OR_9)-R_4$ , or  $-X-C(=N-O-R_8)-R_4$  group at the  $R_1$  position using the methods described in International Publication Nos. WO2005/051317 (Krepeski *et al.*) and WO2005/051324 (Krepeski *et al.*). Compounds of Formula VI prepared from amino esters can also be used to prepare compounds of Formula I having an  $-X-C(R_6)-N(R_8)-R_4$  group at the  $R_1$  position using conventional acyl transfer reaction conditions. Amino alcohols of Formula  $H_2N-X-OH$  can be used to prepare compounds of Formula VI, which can be converted in subsequent steps to a compound of Formula I having an  $-X-S(O)_{0-2}-R_4$ ,  $-X-S(O)_{0-2}-N(R_8)-R_4$ ,  $-X-O-N(R_8)-Q-R_4$ ,  $-X-O-N=C(R_4)-R_4$ ,  $-X-CH(-N(-O-R_8)-Q-R_4)-R_4$  group using methods described in U. S. Patent No. 6,797,718 (Dellaria *et al.*) and International Publication Nos. WO2005/066169 (Bonk and Dellaria), WO2005/018551 (Kshirsagar *et al.*), WO2005/018556 (Kshirsagar *et al.*), and WO2005/051324 (Krepeski *et al.*), respectively.

The amine used in step (1) may be *tert*-butyl carbazate, and the resulting *tert*-butyl 2-(2-chloro-3-nitropyridin-4-yl)hydrazinecarboxylate can be subjected to the conditions of steps (2) to (4). The compound of Formula IX wherein  $R_1$  is a Boc-protected amino group can be deprotected to provide a 1-amino compound or a salt (for example, hydrochloride salt) thereof. The deprotection can be carried out by heating at reflux a solution of a compound of Formula IX in ethanolic hydrogen chloride. The resulting compound of

Formula IX wherein  $R_1$  is an amino group can be treated with a ketone, aldehyde, or corresponding ketal or acetal thereof, under acidic conditions. For example, a ketone can be added to a solution of the hydrochloride salt of a compound of Formula IX in which  $R_1$  is an amino group in a suitable solvent such as isopropanol or acetonitrile in the presence of an acid such as pyridinium *p*-toluene sulfonate or acetic acid, or an acid resin, for example, DOWEX W50-X1 acid resin. The reaction can be performed at an elevated temperature. The resulting imine can be reduced to provide a compound of Formula IX in which  $R_1$  is  $-N(R_1')-Q-R_4$ , wherein  $Q$  is a bond. The reduction can be carried out at room temperature with sodium borohydride in a suitable solvent, for example, methanol. The deprotection shown in step (5) can then be carried out to provide a compound of Formula I. A *tert*-butyl 2-(2-chloro-3-nitropyridin-4-yl)hydrazinecarboxylate of Formula VI can also be manipulated in subsequent steps using the methods described in International Publication No. WO2006/026760 (Stoermer *et al.*) to provide other compounds of Formula I, wherein  $R_1$  is  $-N(R_1')-Q-R_4$ ,  $-N(R_1')-X_1-Y_1-R_4$ , or  $-N(R_1')-X_1-R_{5b}$ .


In certain preferred embodiments,  $R_1$  is a (3-substituted-isoxazol-5-yl)methyl group. This group can be prepared by using propargyl amine as the amine of Formula  $R_1-NH_2$  in step (1) of Reaction Scheme I to provide a compound of Formula VI wherein  $R_1$  is  $-CH_2-C\equiv CH$ . Prior to step (5) of Reaction Scheme I, the alkyne group at the  $R_1$  position undergoes a cycloaddition reaction with a nitrile oxide formed from an  $\alpha$ -chloroaldoxime to provide an isoxazole-substituted 1*H*-imidazo[4,5-*c*]pyridin-2-ol of Formula I.  $\alpha$ -Chloroaldoximes can be prepared by treating an aldoxime with *N*-chlorosuccinimide in a suitable solvent such as DMF. The reaction may be carried out initially below room temperature, at 0 °C for example, and then heated at an elevated temperature in the range of 40 °C to 50 °C. Aldoximes are commercially available or can be prepared from aldehydes by methods well known to one skilled in the art. The resulting  $\alpha$ -chloroaldoxime can optionally be isolated before it is combined with a compound of Formula IX, wherein  $R_1$  is  $-CH_2-C\equiv CH$ , in the presence of a base such as triethylamine to generate a nitrile oxide *in situ* and effect the cycloaddition reaction. The reaction with an  $\alpha$ -chloroaldoxime can be carried out at room temperature in a suitable solvent such as dichloromethane. Other amines of Formula  $NH_2-X-CH=CH_2$  or

NH<sub>2</sub>-X-C≡C-H can also be used in step (1) of Reaction Scheme I to provide compounds of Formula I wherein R<sub>1</sub> is a (3-substituted-isoxazol-5-yl)alkyl group or a (3-substituted-4,5-dihydroisoxazol-5-yl)alkyl group.

Synthetic elaboration can also be carried out at the R<sub>A</sub> or R<sub>B</sub> position of a compound of Formula I or an intermediate of Formula V through IX. For example, the compounds of Formula V in which R<sub>B</sub> is a methyl group are known and can be treated according to steps (1) through (4) of Reaction Scheme I to prepare protected 1*H*-imidazo[4,5-*c*]pyridin-2-ols of Formula IX. The methyl group at the R<sub>B</sub> position can then be brominated using *N*-bromosuccinimide according to the method of Rama Rao, A. V. *et al. Tetrahedron Lett.*, 34, p. 2665, (1993) or Clive, D. L. J. *et al. J. Am. Chem. Soc.*, 116, p. 11275, (1994). The bromine can then be displaced with a variety of primary amines or alkoxide groups using conventional methods, and subsequently step (5) can be followed to provide compounds of Formula I in which R<sub>B</sub> is a methyl group that is substituted by alkylamino, alkoxy, aryloxy, arylalkyleneoxy, heteroaryloxy, or heteroarylalkyleneoxy.


In another example, the compounds of Formula IX wherein R<sub>B</sub> is chloro or bromo, which can be prepared using the methods described in Rousseau, R. J., Robins, R. K., *J. Heterocycl. Chem.*, 2, 196 (1965), may be converted to the corresponding compounds wherein R<sub>B</sub> is alkylamino through palladium-catalyzed coupling with various amines (Wagaw, S., Buchwald, S. L., *J. Org. Chem.*, 61, 7240, (1996)). Likewise, the corresponding compounds wherein R<sub>B</sub> is alkoxy may be prepared by palladium-catalyzed coupling with the desired alcohol (Palucki, M., Wolfe, J. P., Buchwald, S. L., *J. Am. Chem. Soc.*, 119, 3395, (1997)). Displacement of the 6-chloro group with an alkoxide anion could also provide the corresponding 6-alkoxy derivatives (Japanese Patent No. 04018073 (Tenma *et al.*)).

The compounds of Formula IX wherein R<sub>B</sub> is chloro and R<sub>A</sub> is hydrogen may alternatively be prepared according to Reaction Scheme I, using the compounds of Formula VII wherein R<sub>B</sub> is chloro and R<sub>A</sub> is hydrogen. Such compounds of Formula VII can be accessed by reacting a substituted 2,6-dichloro-3-nitropyridin-4-amine of the



formula with an amine of the formula (Bn)<sub>2</sub>NH. The substituted 2,6-dichloro-3-nitropyridin-4-amine can be prepared by reacting 2,6-dichloro-3-nitropyridin-

4-amine with a halogen substituted compound of the formula  $R_1X$  in the presence of a base, such as triethylamine. The 2,6-dichloro-3-nitropyridin-4-amine can be prepared by nitrating 2,6-dichloropyridin-4-amine in the presence of concentrated sulfuric acid/nitric acid (10/90) at a reduced temperature, for example, at 0 °C to form 2,6-dichloro-4-

5 nitraminopyridine, , which can be converted to 2,6-dichloro-3-nitropyridin-4-amine in the presence of concentrated sulfuric acid at an elevated temperature, such as heating over a steam bath (Rousseau, R. J., Robins, R. K., *J. Heterocycl. Chem.*, 2, 196 (1965)). The compounds of Formula IX wherein  $R_B$  is chloro may then be converted to the corresponding compounds wherein  $R_B$  is alkylamino or alkoxy as described above.

Alternatively, compounds of Formula IX wherein  $R_B$  is alkoxy can be prepared according to Reaction Scheme I using compounds of Formula VII wherein  $R_B$  is chloro and  $R_A$  and  $R_1$  are both hydrogen. The compounds of Formula VII wherein  $R_B$  is chloro can be converted to the corresponding compounds wherein  $R_B$  is alkoxy by displacing the chloro group using a metal alkoxide, such as a sodium alkoxide. The reaction can be carried out by adding a 4-amino-6-chloro-3-nitropyridine of Formula VII, where  $R_B$  is chloro and  $R_A$  and  $R_1$  are both hydrogen, in a suitable solvent, such as tetrahydrofuran, to a metal alkoxide solution at a reduced temperature, such as at ice bath temperature, and then heating at an elevated temperature, for example, at 85 °C after completing the addition. In step (3) of Reaction Scheme I, the resulting 4-amino-6-alkoxy-3-nitropyridine of Formula VII, where  $R_B$  is alkoxy and  $R_A$  and  $R_1$  are both hydrogen, can then be reduced to a 3,4-diamino-6-alkoxypyridine of Formula VIII, where  $R_B$  is alkoxy and  $R_A$  and  $R_1$  are both hydrogen. The reduction is conveniently carried out by adding aqueous sodium hydrosulfite to a 4-amino-6-alkoxy-3-nitropyridine of Formula VII in a suitable solvent or solvent mixture such as ethanol/acetonitrile. The reaction can be carried out at room temperature. In step (4) of Reaction Scheme I, a 3,4-diamino-6-alkoxypyridine of Formula VIII can be cyclized to provide a 6-alkoxy-1,3-dihydroimidazo[4,5-*c*]pyridin-2-one, which is the keto tautomer of Formula IX, where  $R_1$  is hydrogen. The cyclization can be conveniently carried out by heating a 3,4-diamino-6-alkoxypyridine of Formula VIII with 1,1'-carbonyldiimidazole in a suitable solvent such as tetrahydrofuran (THF), *tert*-

butyl methyl ether, dichloromethane, or DMF. The reaction may be carried out at room temperature or, preferably, at an elevated temperature such as the reflux temperature of the solvent. The 1-position of the keto tautomer of Formula IX can be substituted by reaction with a compound of the formula X-R<sub>1</sub> wherein X is a halogen, such as a bromo group, and R<sub>1</sub> is other than hydrogen. The reaction can be carried out by heating a keto tautomer of Formula IX with a compound of formula X-R<sub>1</sub> in a suitable solvent, such as DMF, at an elevated temperature, for example, 80 °C to form a compound of Formula IX, substituted at the 1-position with R<sub>1</sub>.

Likewise, compounds of Formula IX wherein R<sub>B</sub> is alkylamino can be prepared according Scheme I using compounds of Formula VII wherein R<sub>B</sub> is chloro and R<sub>A</sub> and R<sub>1</sub> are both hydrogen. The compounds of Formula VII wherein R<sub>B</sub> is chloro can be converted to the corresponding compounds wherein R<sub>B</sub> is alkylmino by displacing the chloro group using an excess, such as five equivalents, of an alkylamine, such as, for example, *n*-butylamine. The reaction can be carried out by adding a 4-amino-6-chloro-3-nitropyridine of Formula VII, where R<sub>B</sub> is chloro and R<sub>A</sub> and R<sub>1</sub> are both hydrogen, in a suitable solvent, such as trifluoroethanol, to a solution of the desired alkylamine and then heating at an elevated temperature, for example, at 130 °C in a sealed tube for a period of time, for example, eighteen to twenty-four hours. Steps (3) and (4) of Reaction Scheme I and the installation of an R<sub>1</sub> group other than hydrogen can then be carried out as described above to provide a 1-substituted, 6-alkylamino compound of Formula IX.


In a further alternative, compounds of Formula IX wherein R<sub>B</sub> is alkoxy may be accessed by O-alkylation of the corresponding 6-oxo-1(3)H-imidazo[4,5-c]pyridin-4-ylamine by O-alkylation methods utilizing a base such as cesium carbonate in a solvent such as DMF (Meurer, L. *et al.*, *Bioorg. Med. Chem. Lett.*, 15(3) 645, (2005)). In addition, O-alkylation could be accomplished under Mitsunobu conditions (Li, Q. *et al.*, *Bioorg. Med. Chem. Lett.*, 16(6), 1679 (2006)).

Step (6) of Reaction Scheme I can be used to prepare a compound of Formula II. The amino group of a pyridine of Formula I can be converted by conventional methods to a functional group such as an amide, carbamate, urea, amidine, or another hydrolyzable group. A compound of this type can be made by the replacement of a hydrogen atom in an amino group with a group such as -C(O)-R',  $\alpha$ -aminoacyl,  $\alpha$ -aminoacyl- $\alpha$ -aminoacyl, -C(O)-O-R', -C(O)-N(R'')-R', -C(=NY')-R', -CH(OH)-C(O)-OY', -CH(OC<sub>1-4</sub> alkyl)Y<sub>0</sub>,

-CH<sub>2</sub>Y<sub>1</sub>, or -CH(CH<sub>3</sub>)Y<sub>1</sub>; wherein R' and R'' are each independently C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, phenyl, or benzyl, each of which may be unsubstituted or substituted by one or more substituents independently selected from the group consisting of halogen, hydroxy, nitro, cyano, carboxy, C<sub>1-6</sub> alkyl, C<sub>1-4</sub> alkoxy, aryl, heteroaryl, arylC<sub>1-4</sub> alkylenyl, heteroarylC<sub>1-4</sub> alkylenyl, haloC<sub>1-4</sub> alkylenyl, haloC<sub>1-4</sub> alkoxy, -O-C(O)-CH<sub>3</sub>, -C(O)-O-CH<sub>3</sub>, -C(O)-NH<sub>2</sub>, -O-CH<sub>2</sub>-C(O)-NH<sub>2</sub>, -NH<sub>2</sub>, and -S(O)<sub>2</sub>-NH<sub>2</sub>; with the proviso that R'' may also be hydrogen; each  $\alpha$ -aminoacyl group is independently selected from racemic, D, or L-amino acids; Y' is hydrogen, C<sub>1-6</sub> alkyl, or benzyl; Y<sub>0</sub> is C<sub>1-6</sub> alkyl, carboxyC<sub>1-6</sub> alkylenyl, aminoC<sub>1-4</sub> alkylenyl, mono-*N*-C<sub>1-6</sub> alkylaminoC<sub>1-4</sub> alkylenyl, or di-*N,N*-C<sub>1-6</sub> alkylaminoC<sub>1-4</sub> alkylenyl; and Y<sub>1</sub> is mono-*N*-C<sub>1-6</sub> alkylamino, di-*N,N*-C<sub>1-6</sub> alkylamino, morpholin-4-yl, piperidin-1-yl, pyrrolidin-1-yl, or 4-C<sub>1-4</sub> alkylpiperazin-1-yl. Particularly useful compounds of Formula II are amides derived from carboxylic acids containing one to ten carbon atoms, amides derived from amino acids, and carbamates containing one to ten carbon atoms. The reaction can be carried out, for example, by combining a compound of Formula I with a chloroformate or acid chloride, such as ethyl chloroformate or acetyl chloride, in the presence of a base such as triethylamine in a suitable solvent such as dichloromethane at room temperature.

Step (6a) of Reaction Scheme I can be used to prepare a compound of Formula III. The hydrogen atom of the alcohol group of Formula I can be replaced using conventional methods with a group such as C<sub>1-6</sub> alkanoyloxymethyl, 1-(C<sub>1-6</sub> alkanoyloxy)ethyl, 1-methyl-1-(C<sub>1-6</sub> alkanoyloxy)ethyl, C<sub>1-6</sub> alkoxycarbonyloxymethyl, *N*-(C<sub>1-6</sub> alkoxycarbonyl)aminomethyl, succinoyl, C<sub>1-6</sub> alkanoyl,  $\alpha$ -aminoC<sub>1-4</sub> alkanoyl, arylacyl, -P(O)(OH)<sub>2</sub>, -P(O)(O-C<sub>1-6</sub> alkyl)<sub>2</sub>, C<sub>1-6</sub> alkoxycarbonyl, C<sub>1-6</sub> alkylcarbamoyl, and  $\alpha$ -aminoacyl or  $\alpha$ -aminoacyl- $\alpha$ -aminoacyl, where each  $\alpha$ -aminoacyl group is independently selected from racemic, D, and L-amino acids. Particularly useful compounds of Formula III are esters made from carboxylic acids containing one to six carbon atoms, unsubstituted or substituted benzoic acid esters, or esters made from naturally occurring amino acids.

## Reaction Scheme I



Certain compounds of Formula V can be prepared according to Reaction Scheme II, wherein  $R_{11}$  and Boc are as defined above,  $R_{Bx}$  is alkenyl,  $-R_{11}$ , or a carboxy group, and  $R_{Ba}$  is alkenyl,  $-R_{11}$  or  $-NHR_{11}$ . A 4-hydroxy-2H-pyran-2-one of Formula X in which  $R_{Bx}$  is alkenyl or  $-R_{11}$  can be prepared from  $\beta,\gamma$ -diketoesters according to the method of Lygo, B., *Tetrahedron*, 51, pp.12859-12868, (1995) or Song, D. *et al.*, *Tetrahedron*, 59, pp. 6899-6904, (2003). The compound of Formula X in which  $R_{Bx}$  is methyl is commercially available and can undergo lithiation-substitution reactions using the method of Poulton, G. A., and Cyr, T. D., *Can. J. Chem.* 58, p. 2158, (1980) to provide compounds of Formula X in which  $R_{Bx}$  is  $-R_{11}$ . The compound of Formula X in which  $R_{Bx}$  is a carboxy group can be prepared by the method of Stetter, H. and Schellhammer, C.-W., *Chem. Ber.*, 90, p. 755 (1957).



In step (1) of Reaction Scheme II, a 4-hydroxy-2*H*-pyran-2-one of Formula X is converted to a pyridin-2,4-diol of Formula XI. The reaction can be carried out by heating a compound of Formula X in aqueous ammonium hydroxide at a temperature of 80 °C to 130 °C, preferably at a temperature of about 100 °C to about 120 °C.


5 In step (2) of Reaction Scheme II, a compound of Formula XI in which R<sub>Bx</sub> is a carboxy group is treated with diphenylphosphoryl azide to provide an azide of Formula XII, which undergoes a Curtius rearrangement in step (3) to provide a carbamate-substituted pyridin-2,4-diol of Formula XIII. The Curtius rearrangement in step (3) can be carried out by heating at an elevated temperature such as 70 °C to 110 °C in a suitable  
10 solvent such as *tert*-butanol to provide the *tert*-butyl carbamate of Formula XIII.

In step (4) of Reaction Scheme II, a carbamate-substituted pyridin-2,4-diol of Formula XIII is deprotected using conventional methods. For example, the Boc group can be removed by treating with trifluoroacetic acid at room temperature to provide an amino-substituted pyridin-2,4-diol of Formula XIV.

15 In step (5) of Reaction Scheme II, an amino-substituted pyridin-2,4-diol of Formula XIV reacts with an aldehyde or ketone to provide an imine. Numerous aldehydes and ketones are commercially available; others can be readily prepared using known synthetic methods. The reaction can be conveniently carried out by combining the aldehyde or ketone with a compound of Formula XIV in a suitable solvent such as  
20 methanol. The reaction can be carried out at room temperature, or at an elevated temperature. Optionally, an acid such as pyridine hydrochloride can be added. The imine is then reduced to provide an amino-substituted of Formula pyridin-2,4-diol of Formula XV. The reduction is conveniently carried out by treating the oxime with excess sodium cyanoborohydride in a suitable solvent or solvent mixture such as methanol/acetic acid.  
25 Optionally, hydrochloric acid may be added. The reaction can be carried out at room temperature or at an elevated temperature.

In steps (6) and (7) of Reaction Scheme II, a compound of Formula XI or XV is converted to a 2,4-dichloro-3-nitropyridine of Formula Va by treating first with nitric acid and then with phosphorus(III) oxychloride according to known methods. See, for  
30 example, the methods in U.S. Patent Nos. 5,446,153 (Lindstrom *et al.*) and 6,525,064 (Dellaria *et al.*).

## Reaction Scheme II



Compounds of the invention can also be prepared using variations of the synthetic routes shown in Reaction Schemes I and II that would be apparent to one of skill in the art, including variations described in the EXAMPLES below.

## Pharmaceutical Compositions and Biological Activity

Pharmaceutical compositions of the invention contain a therapeutically effective amount of a compound or salt of the invention as described above in combination with a pharmaceutically acceptable carrier.

The terms "a therapeutically effective amount" and "effective amount" mean an amount of the compound or salt sufficient to induce a therapeutic or prophylactic effect, such as cytokine induction, cytokine inhibition, immunomodulation, antitumor activity, and/or antiviral activity. The exact amount of compound or salt used in a pharmaceutical composition of the invention will vary according to factors known to those of skill in the art, such as the physical and chemical nature of the compound or salt, the nature of the carrier, and the intended dosing regimen.

In some embodiments, the compositions of the invention will contain sufficient active ingredient or prodrug to provide a dose of about 100 nanograms per kilogram (ng/kg) to about 50 milligrams per kilogram (mg/kg), preferably about 10 micrograms per kilogram ( $\mu\text{g/kg}$ ) to about 5 mg/kg, of the compound or salt to the subject.

In other embodiments, the compositions of the invention will contain sufficient active ingredient or prodrug to provide a dose of, for example, from about 0.01 mg/m<sup>2</sup> to

about 5.0 mg/m<sup>2</sup>, computed according to the Dubois method, in which the body surface area of a subject (m<sup>2</sup>) is computed using the subject's body weight:  $m^2 = (wt\ kg^{0.425} \times height\ cm^{0.725}) \times 0.007184$ , although in some embodiments the methods may be performed by administering a compound or salt or composition in a dose outside this range. In some of these embodiments, the method includes administering sufficient compound to provide a dose of from about 0.1 mg/m<sup>2</sup> to about 2.0 mg/m<sup>2</sup> to the subject, for example, a dose of from about 0.4 mg/m<sup>2</sup> to about 1.2 mg/m<sup>2</sup>.

A variety of dosage forms may be used, such as tablets, lozenges, capsules, parenteral formulations, syrups, creams, ointments, aerosol formulations, transdermal patches, transmucosal patches and the like. These dosage forms can be prepared with conventional pharmaceutically acceptable carriers and additives using conventional methods, which generally include the step of bringing the active ingredient into association with the carrier.

The compounds or salts of the invention can be administered as the single therapeutic agent in the treatment regimen, or the compounds or salts described herein may be administered in combination with one another or with other active agents, including additional immune response modifiers, antivirals, antibiotics, antibodies, proteins, peptides, oligonucleotides, etc.

Compounds or salts of the invention have been shown to induce the production of certain cytokines in experiments performed according to the tests set forth below. These results indicate that the compounds or salts are useful for modulating the immune response in a number of different ways, rendering them useful in the treatment of a variety of disorders.

In some embodiments, compounds or salts of Formula I can be especially useful as immune response modifiers due to their ability to selectively induce IFN- $\alpha$ . As used herein, to "selectively induce IFN- $\alpha$ " means, that when tested according to the test methods described herein, the effective minimum concentration (of the compound or salt) for IFN- $\alpha$  induction is less than the effective minimum concentration for TNF- $\alpha$  induction. In some embodiments, the effective minimum concentration for IFN- $\alpha$  induction is at least 3-fold less than the effective minimum concentration for TNF- $\alpha$  induction. In some embodiments, the effective minimum concentration for IFN- $\alpha$  induction is at least 6-fold less than the effective minimum concentration for TNF- $\alpha$  induction. In other

embodiments, the effective minimum concentration for IFN- $\alpha$  induction is at least 10-fold less than the effective minimum concentration for TNF- $\alpha$  induction. In other embodiments, the effective minimum concentration for IFN- $\alpha$  induction is at least 100-fold less than the effective minimum concentration for TNF- $\alpha$  induction. In some  
5      embodiments, when tested according to the test methods described herein, the amount TNF- $\alpha$  induced by compounds of the invention is at or below the background level of TNF- $\alpha$  in the test method. Compounds or salts of the invention may, therefore, provide a benefit, for example, a reduced inflammatory response, particularly when administered systemically, over compounds that also induce pro-inflammatory cytokines (e.g. TNF- $\alpha$ )  
10     or that induce pro-inflammatory cytokines at higher levels.

Cytokines whose production may be induced by the administration of compounds or salts of the invention generally include interferon- $\alpha$  (IFN- $\alpha$ ) and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) as well as certain interleukins (IL). Cytokines whose biosynthesis may be induced by compounds or salts of the invention include IFN- $\alpha$ , TNF- $\alpha$ , IL-1, IL-6, IL-10  
15     and IL-12, and a variety of other cytokines. Among other effects, these and other cytokines can inhibit virus production and tumor cell growth, making the compounds or salts useful in the treatment of viral diseases and neoplastic diseases. Accordingly, the invention provides a method of inducing cytokine biosynthesis in an animal comprising administering an effective amount of a compound or salt or composition of the invention  
20     to the animal. The animal to which the compound or salt or composition is administered for induction of cytokine biosynthesis may have a disease as described *infra*, for example a viral disease or a neoplastic disease, and administration of the compound or salt may provide therapeutic treatment. Alternatively, the compound or salt may be administered to the animal prior to the animal acquiring the disease so that administration of the  
25     compound or salt may provide a prophylactic treatment.

In addition to the ability to induce the production of cytokines, compounds or salts described herein can affect other aspects of the innate immune response. For example, natural killer cell activity may be stimulated, an effect that may be due to cytokine induction. The compounds or salts may also activate macrophages, which in turn  
30     stimulate secretion of nitric oxide and the production of additional cytokines. Further, the compounds or salts may cause proliferation and differentiation of B-lymphocytes.

Compounds or salts of the invention can also have an effect on the acquired immune response. For example, the production of the T helper type 1 (T<sub>H</sub>1) cytokine IFN- $\gamma$  may be induced indirectly and the production of the T helper type 2 (T<sub>H</sub>2) cytokines IL-4, IL-5 and IL-13 may be inhibited upon administration of the compounds or salts.

5 Whether for prophylaxis or therapeutic treatment of a disease, and whether for effecting innate or acquired immunity, the compound or salt or composition may be administered alone or in combination with one or more active components as in, for example, a vaccine adjuvant. When administered with other components, the compound or salt and other component or components may be administered separately; together but  
10 independently such as in a solution; or together and associated with one another such as (a) covalently linked or (b) non-covalently associated, e.g., in a colloidal suspension.

Conditions for which compounds or salts identified herein may be used as treatments include, but are not limited to:

(a) viral diseases such as, for example, diseases resulting from infection by an  
15 adenovirus, a herpesvirus (e.g., HSV-I, HSV-II, CMV, or VZV), a poxvirus (e.g., an orthopoxvirus such as variola or vaccinia, or molluscum contagiosum), a picornavirus (e.g., rhinovirus or enterovirus), an orthomyxovirus (e.g., influenzavirus), a paramyxovirus (e.g., parainfluenzavirus, mumps virus, measles virus, and respiratory syncytial virus (RSV)), a coronavirus (e.g., SARS), a papovavirus (e.g., papillomaviruses, such as those  
20 that cause genital warts, common warts, or plantar warts), a hepadnavirus (e.g., hepatitis B virus), a flavivirus (e.g., hepatitis C virus or Dengue virus), or a retrovirus (e.g., a lentivirus such as HIV);

(b) bacterial diseases such as, for example, diseases resulting from infection by bacteria of, for example, the genus Escherichia, Enterobacter, Salmonella, Staphylococcus,  
25 Shigella, Listeria, Aerobacter, Helicobacter, Klebsiella, Proteus, Pseudomonas, Streptococcus, Chlamydia, Mycoplasma, Pneumococcus, Neisseria, Clostridium, Bacillus, Corynebacterium, Mycobacterium, Campylobacter, Vibrio, Serratia, Providencia, Chromobacterium, Brucella, Yersinia, Haemophilus, or Bordetella;

(c) other infectious diseases, such chlamydia, fungal diseases including but not  
30 limited to candidiasis, aspergillosis, histoplasmosis, cryptococcal meningitis, or parasitic diseases including but not limited to malaria, pneumocystis carinii pneumonia, leishmaniasis, cryptosporidiosis, toxoplasmosis, and trypanosome infection;

(d) neoplastic diseases, such as intraepithelial neoplasias, cervical dysplasia, actinic keratosis, basal cell carcinoma, squamous cell carcinoma, renal cell carcinoma, Kaposi's sarcoma, melanoma, leukemias including but not limited to acute myeloid leukemia, acute lymphocytic leukemia, chronic myeloid leukemia, chronic lymphocytic leukemia, multiple myeloma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cutaneous T-cell lymphoma, B-cell lymphoma, and hairy cell leukemia, and other cancers;

(e)  $T_H2$ -mediated, atopic diseases, such as atopic dermatitis or eczema, eosinophilia, asthma, allergy, allergic rhinitis, and Ommen's syndrome;

(f) certain autoimmune diseases such as systemic lupus erythematosus, essential thrombocythaemia, multiple sclerosis, discoid lupus, alopecia areata; and

(g) diseases associated with wound repair such as, for example, inhibition of keloid formation and other types of scarring (e.g., enhancing wound healing, including chronic wounds).

Additionally, a compound or salt of the present invention may be useful as a vaccine adjuvant for use in conjunction with any material that raises either humoral and/or cell mediated immune response, such as, for example, live viral, bacterial, or parasitic immunogens; inactivated viral, tumor-derived, protozoal, organism-derived, fungal, or bacterial immunogens; toxoids; toxins; self-antigens; polysaccharides; proteins; glycoproteins; peptides; cellular vaccines; DNA vaccines; autologous vaccines; recombinant proteins; and the like, for use in connection with, for example, BCG, cholera, plague, typhoid, hepatitis A, hepatitis B, hepatitis C, influenza A, influenza B, parainfluenza, polio, rabies, measles, mumps, rubella, yellow fever, tetanus, diphtheria, hemophilus influenza b, tuberculosis, meningococcal and pneumococcal vaccines, adenovirus, HIV, chicken pox, cytomegalovirus, dengue, feline leukemia, fowl plague, HSV-1 and HSV-2, hog cholera, Japanese encephalitis, respiratory syncytial virus, rotavirus, papilloma virus, yellow fever, and Alzheimer's Disease.

Compounds or salts of the present invention may be particularly helpful in individuals having compromised immune function. For example, compounds or salts may be used for treating the opportunistic infections and tumors that occur after suppression of cell mediated immunity in, for example, transplant patients, cancer patients and HIV patients.

Thus, one or more of the above diseases or types of diseases, for example, a viral disease or a neoplastic disease may be treated in an animal in need thereof (having the disease) by administering a therapeutically effective amount of a compound or salt of Formula I, II, III, any of the embodiments described herein, or a combination thereof to the animal.

An animal may also be vaccinated by administering an effective amount of a compound or salt of Formula I, II, III, any of the embodiments described herein, or a combination thereof to the animal as a vaccine adjuvant. In one embodiment, there is provided a method of vaccinating an animal comprising administering an effective amount of a compound or salt described herein to the animal as a vaccine adjuvant.

An amount of a compound or salt effective to induce cytokine biosynthesis is an amount sufficient to cause one or more cell types, such as monocytes, macrophages, dendritic cells and B-cells to produce an amount of one or more cytokines such as, for example, IFN- $\alpha$ , TNF- $\alpha$ , IL-1, IL-6, IL-10 and IL-12 that is increased (induced) over a background level of such cytokines. The precise amount will vary according to factors known in the art but is expected to be a dose of about 100 ng/kg to about 50 mg/kg, preferably about 10  $\mu$ g/kg to about 5 mg/kg. In other embodiments, the amount is expected to be a dose of, for example, from about 0.01 mg/m<sup>2</sup> to about 5.0 mg/m<sup>2</sup>, (computed according to the Dubois method as described above) although in some embodiments the induction or inhibition of cytokine biosynthesis may be performed by administering a compound or salt in a dose outside this range. In some of these embodiments, the method includes administering sufficient compound or salt or composition to provide a dose of from about 0.1 mg/m<sup>2</sup> to about 2.0 mg/m<sup>2</sup> to the subject, for example, a dose of from about 0.4 mg/m<sup>2</sup> to about 1.2 mg/m<sup>2</sup>.

The invention also provides a method of treating a viral infection in an animal and a method of treating a neoplastic disease in an animal comprising administering an effective amount of a compound or salt or composition of the invention to the animal. An amount effective to treat or inhibit a viral infection is an amount that will cause a reduction in one or more of the manifestations of viral infection, such as viral lesions, viral load, rate of virus production, and mortality as compared to untreated control animals. The precise amount that is effective for such treatment will vary according to factors known in the art but is expected to be a dose of about 100 ng/kg to about 50 mg/kg,

preferably about 10 µg/kg to about 5 mg/kg. An amount of a compound or salt effective to treat a neoplastic condition is an amount that will cause a reduction in tumor size or in the number of tumor foci. Again, the precise amount will vary according to factors known in the art but is expected to be a dose of about 100 ng/kg to about 50 mg/kg, preferably  
5 about 10 µg/kg to about 5 mg/kg. In other embodiments, the amount is expected to be a dose of, for example, from about 0.01 mg/m<sup>2</sup> to about 5.0 mg/m<sup>2</sup>, (computed according to the Dubois method as described above) although in some embodiments either of these methods may be performed by administering a compound or salt in a dose outside this range. In some of these embodiments, the method includes administering sufficient  
10 compound or salt to provide a dose of from about 0.1 mg/m<sup>2</sup> to about 2.0 mg/m<sup>2</sup> to the subject, for example, a dose of from about 0.4 mg/m<sup>2</sup> to about 1.2 mg/m<sup>2</sup>.

The methods of the invention may be performed on any suitable subject. Suitable subjects include but are not limited to animals such as but not limited to humans, non-human primates, rodents, dogs, cats, horses, pigs, sheep, goats, or cows.

15 In addition to the formulations and uses described specifically herein, other formulations, uses, and administration devices suitable for compounds of the present invention are described in, for example, International Publication Nos. WO 03/077944 and WO 02/036592, U.S. Patent No. 6,245,776, and U.S. Publication Nos. 2003/0139364, 2003/185835, 2004/0258698, 2004/0265351, 2004/076633, and 2005/0009858.

20

Objects and advantages of this invention are further illustrated by the following examples, but the particular materials and amounts thereof recited in these examples, as well as other conditions and details, should not be construed to unduly limit this invention.

25

## EXAMPLES

In the examples below automated flash chromatography was carried out using a HORIZON HPFC system (an automated high-performance flash purification product available from Biotage, Inc, Charlottesville, Virginia, USA). For some of these purifications, either a FLASH 40+M silica cartridge or a FLASH 25+M silica cartridge  
30 (both available from Biotage, Inc, Charlottesville, Virginia, USA) was used. In some chromatographic separations, the solvent mixture 80/18/2 v/v/v



chloroform/methanol/concentrated ammonium hydroxide (CMA) was used as the polar component of the eluent. In these separations, CMA was mixed with chloroform in the indicated ratio.

5 Preparation of *N,N*-Bis(4-methoxybenzyl)amine

Part A


4-Methoxybenzylamine (40 g, 290 mmol) was cooled to 0 °C, and *p*-anisaldehyde (39.7 g, 292 mmol) was added dropwise. The reaction was stirred at ambient temperature for two hours, concentrated under reduced pressure, and further dried under high vacuum  
10 overnight to provide 97 g of *N*-(4-methoxybenzyl)-*N*-[(4-methoxyphenyl)methylidene]amine as a white, waxy solid.

Part B

A solution of the material from Part A in ethanol (300 mL) was cooled to 0 °C and  
15 stirred rapidly. Solid sodium borohydride (22.1 g, 584 mmol) was added slowly over a period of several minutes, and the reaction was stirred at ambient temperature for two hours. Water (300 mL) was added, and the resulting mixture was shaken and allowed to stand overnight. The mixture was extracted with diethyl ether (3 x 100 mL), and the combined extracts were washed with water (200 mL), dried over magnesium sulfate,  
20 filtered through a layer of CELITE filter agent, concentrated under reduced pressure, and further dried under high vacuum to provide 67 g of *N,N*-bis(4-methoxybenzyl)amine as a white solid.

Example 1

4-Amino-1-benzyl-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol



25

Part A

A solution of hexanoic acid (2.51 mL, 0.0200 mol) in dichloromethane (5 mL) was cooled to 0 °C, and oxalyl chloride (8.7 mL, 0.10 mol) was added. The solution was

allowed to warm to room temperature and stirred for 20 hours under a nitrogen atmosphere. The solvent was removed under reduced pressure, and the residue was dissolved in hexane (50 mL). The solution was cooled to 0 °C, and triethylamine (3.1 mL, 22 mmol) and 2-methylaziridine (1.57 mL of 90% pure material, 20 mmol) were sequentially added. The resultant mixture was stirred for one hour under a nitrogen atmosphere, diluted with ethyl acetate (50 mL), and filtered through a layer of CELITE filter agent. The filtrate was concentrated under reduced pressure, and the residue was purified by flash chromatography on silica gel (eluting with 11% ethyl acetate in hexane) to provide 1.91 g of 1-hexanoyl-2-methylaziridine as a yellow oil.

10 Part B

Sodium hydride (537 mg of a 60% dispersion in mineral oil, 13.4 mmol) was washed three times with hexanes and then suspended in tetrahydrofuran (THF) (30 mL). A solution of *tert*-butyl acetoacetate (1.94 g, 12.2 mmol) in THF (10 mL) was added dropwise to the suspension, and the mixture was stirred for 30 minutes and then cooled to 0 °C. *n*-Butyllithium (8.4 mL of a 1.6 M solution in hexane) was added, and the resulting yellow-orange solution was stirred at 0 °C for 20 minutes. A solution of 1-hexanoyl-2-methylaziridine (1.90 g, 12.2 mmol) in THF (10 mL) was added, and the reaction was stirred at 0 °C for 1.5 hours. Saturated aqueous ammonium chloride was added, and the mixture was extracted with ethyl acetate (3 x 40 mL). The combined extracts were dried over magnesium sulfate, filtered, and concentrated under reduced pressure. The crude product was purified by flash chromatography on silica gel (eluting with 5% ethyl acetate in hexane) to provide 1.95 g of *tert*-butyl 3,5-dioxodecanoate as a colorless oil.

20 Part C

Trifluoroacetic acid (16 mL) was added to a solution of *tert*-butyl 3,5-dioxodecanoate (1.95 g, 7.61 mmol) in dichloromethane (45 mL), and the solution was stirred at room temperature for two hours. The volatiles were removed under reduced pressure, and the residue was dissolved in acetic anhydride (44 mL). The solution was stirred overnight at room temperature, and the acetic anhydride was removed under reduced pressure. The residue was dissolved in methanol (30 mL), and potassium carbonate (105 mg, 0.76 mmol) was added. The mixture was stirred for three hours at room temperature, and an analysis by high-performance liquid chromatography (HPLC) indicated the reaction was incomplete. Additional potassium carbonate (100 mg) was

added, and the reaction was stirred for one hour at room temperature. The volatiles were removed under reduced pressure, and the residue was partitioned between saturated aqueous ammonium chloride and dichloromethane. The aqueous layer was separated and extracted with dichloromethane (3 x 50 mL). The combined organic fractions were dried  
5 over magnesium sulfate, filtered, and concentrated under reduced pressure to provide 4-hydroxy-6-pentyl-2*H*-pyran-2-one as an orange oil that solidified upon standing.

#### Part D

A suspension of 4-hydroxy-6-pentyl-2*H*-pyran-2-one (0.750 g, 4.12 mmol) in concentrated aqueous ammonium hydroxide (10 mL) was heated at 100 °C for six hours  
10 and allowed to cool to room temperature. A precipitate was present and was isolated by filtration, triturated with methanol, and isolated by filtration to provide 0.700 g of 4-hydroxy-6-pentylpyridin-2(1*H*)-one as a tan solid.

#### Part E

Fuming nitric acid (20 mL) was carefully added to a suspension of the material  
15 from Part D in water (5 mL), and the reaction was heated at 80 °C for 30 minutes, allowed to cool to room temperature, and poured into ice water. Some of the water was removed under reduced pressure, and a precipitate formed. The mixture was cooled to approximately 0 °C, and the precipitate was collected by filtration and dried under high vacuum to provide 0.620 g of 3-nitro-6-pentylpyridine-2,4-diol as a pale yellow solid.

#### Part F

A solution of 3-nitro-6-pentylpyridine-2,4-diol (1.00 g, 4.42 mmol) in phosphorus(III) oxychloride (15 mL) was heated at 80 °C for four hours. The excess phosphorous(III) oxychloride was removed under reduced pressure, and saturated aqueous sodium bicarbonate was added to adjust the residue to pH 10. The basic mixture was  
25 extracted several times with ethyl acetate, and the combined extracts were washed with brine, dried over magnesium sulfate, filtered through a layer of CELITE filter agent, and concentrated under reduced pressure. The crude product was purified by flash chromatography on silica gel (eluting with 25% ethyl acetate in hexane) to provide 0.780 g of 2,4-dichloro-3-nitro-6-pentylpyridine as a tan oil.

#### Part G

Triethylamine (1.77 mL, 12.7 mmol) and benzylamine (0.83 mL, 7.6 mmol) were added to a solution of 2,4-dichloro-3-nitro-6-pentylpyridine (2.22 g, 8.44 mmol) in *N,N*-

dimethylformamide (DMF) (50 mL), and the solution was stirred overnight at room temperature. The DMF was removed under reduced pressure, and the residue was partitioned between saturated aqueous sodium bicarbonate and dichloromethane. The aqueous layer was separated and extracted with dichloromethane, and the combined  
5 organic fractions were washed with brine, dried over magnesium sulfate, filtered, and concentrated under reduced pressure. The crude product was purified by flash chromatography on silica gel (eluting with 1% to 3% ethyl acetate in hexanes) to provide 1.39 g of *N*-benzyl-2-chloro-3-nitro-6-pentylpyridin-4-amine.

#### Part H

10 Triethylamine (1.44 mL, 10.3 mmol) and *N,N*-bis(4-methoxybenzyl)amine (2.65 g, 10.3 mmol) were added to a solution of *N*-benzyl-2-chloro-3-nitro-6-pentylpyridin-4-amine (2.30 g, 6.89 mmol) in toluene (100 mL), and the yellow solution was heated at reflux overnight under a nitrogen atmosphere. The volatiles were removed under reduced pressure, and the residue was partitioned between ethyl acetate and saturated aqueous  
15 sodium bicarbonate. The aqueous layer was separated and extracted with ethyl acetate, and the combined organic fractions were washed with brine, dried over magnesium sulfate, filtered, and concentrated under reduced pressure to provide *N*<sup>4</sup>-benzyl-*N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-3-nitro-6-pentylpyridine-2,4-diamine, which was used without purification.

#### 20 Part I

Sodium borohydride (0.150 g, 3.97 mmol) was added to a mixture of nickel(II) chloride hexahydrate (0.820 g, 3.45 mmol) in methanol (60 mL), and the mixture was stirred for 15 minutes. A solution of *N*<sup>4</sup>-benzyl-*N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-3-nitro-6-pentylpyridine-2,4-diamine (3.82 g, 6.89 mmol) in methanol (50 mL) and  
25 dichloromethane (25 mL) was added. Additional sodium borohydride (0.319 g, 8.43 mmol) was added in portions over a period of ten minutes, and the mixture was stirred for one hour at room temperature. An analysis by HPLC indicated the presence of starting material, and additional nickel(II) chloride hexahydrate (0.800 g, 3.37 mmol) and sodium borohydride (0.250 g, 6.61 mmol) were added. The reaction was stirred at room  
30 temperature for two hours and then filtered through a layer of CELITE filter agent. The filter cake was washed with dichloromethane, and the filtrate was concentrated under reduced pressure. The residue was partitioned between saturated aqueous sodium

bicarbonate and dichloromethane, and the work-up procedure described in Part G was followed. The crude product was purified by flash chromatography on silica gel (eluting with 2% to 4% methanol in dichloromethane) to provide 3.45 g of *N*<sup>4</sup>-benzyl-*N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-6-pentylpyridine-2,3,4-triamine as a thick, dark oil.

5     Part J

Carbonyl diimidazole (1.60 g, 9.86 mmol) was added to a solution of *N*<sup>4</sup>-benzyl-*N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-6-pentylpyridine-2,3,4-triamine (3.45 g, 6.58 mmol) in THF (50 mL), and the dark green solution was heated at reflux under a nitrogen atmosphere for two hours. The volatiles were removed under reduced pressure, and the residue was  
10     purified by flash chromatography on silica gel (eluting with 2% methanol in dichloromethane) to provide 3.52 g of 1-benzyl-4-[bis(4-methoxybenzyl)amino]-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a thick, yellow oil that solidified upon standing.

Part K


A solution of 1-benzyl-4-[bis(4-methoxybenzyl)amino]-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol (3.52 g, 6.39 mmol) in trifluoroacetic acid (15 mL) was stirred at room  
15     temperature for five hours and then diluted with water. The resulting mixture was adjusted to approximately pH 9 with the addition of solid sodium carbonate. The aqueous layer was separated and extracted several times with dichloromethane and dichloromethane/methanol. The combined organic fractions were dried over magnesium  
20     sulfate, filtered, and concentrated under reduced pressure. The resulting white solid was triturated with acetonitrile/methanol and isolated by filtration to provide 1.08 g of 4-amino-1-benzyl-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a white crystalline solid, mp 260-262 °C.

<sup>1</sup>H NMR (300 MHz, *d*<sub>4</sub>-MeOH) δ 7.57-7.51 (m, 5H), 6.57 (s, 1H), 5.26 (s, 2H), 2.77 (dd, *J*  
25     = 7.4, 7.8 Hz, 2H), 1.82 (m, 2H), 1.53-1.48 (m, 4H), 1.10 (t, *J* = 7.0 Hz, 3H);

MS (APCI) *m/z* 311 (*M* + *H*<sup>+</sup>);

Anal. Calcd for C<sub>18</sub>H<sub>22</sub>N<sub>4</sub>O·0.67 CF<sub>3</sub>CO<sub>2</sub>H: C, 60.06; H, 5.91; N, 14.48. Found: C, 59.72; H, 6.30; N, 14.57.

## Example 2

4-Amino-6-pentyl-1-(2-phenylethyl)-1*H*-imidazo[4,5-*c*]pyridin-2-ol

## Part A

5 Phenethylamine (0.86 mL, 6.8 mmol) was added to a stirred solution of 2,4-dichloro-3-nitro-6-pentylpyridine (2.0 g, 7.6 mmol) and triethylamine (1.6 mL, 11 mmol) in DMF (38 mL), and the reaction was stirred for three hours at room temperature. Water (200 mL) was added, and the mixture was extracted with ethyl acetate (2 x 50 mL). The combined organic fractions were washed with brine (50 mL), dried over sodium sulfate,  
10 filtered, and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel (eluting with 10% ethyl acetate in hexane) to provide 1.1 g of (2-chloro-3-nitro-6-pentylpyridin-4-yl)phenethylamine as a yellow oil.

## Part B

15 Triethylamine (0.66 mL, 4.7 mmol) and *N,N*-bis(4-methoxybenzyl)amine (1.2 g, 4.7 mmol) were added to a solution of (2-chloro-3-nitro-6-pentylpyridin-4-yl)phenethylamine (1.1 g, 3.2 mmol) in toluene (32 mL), and the yellow solution was heated at reflux for three hours, stirred overnight at room temperature, and heated at reflux for two hours. The work-up procedure described in Part H of Example 1 was followed to provide *N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-3-nitro-6-pentyl-*N*<sup>4</sup>-(2-phenylethyl)pyridine-2,4-  
20 diamine as an oil.

## Part C

Sodium borohydride (0.070 g, 1.85 mmol) was added to a mixture of nickel(II) chloride hexahydrate (0.38 g, 1.6 mmol) in methanol (25 mL), and the mixture was stirred for 15 minutes. A solution of the material from Part B in methanol (25 mL) and  
25 dichloromethane (11 mL) was added. Additional sodium borohydride (0.150 g, 3.97 mmol) was added in portions, and the mixture was stirred for one hour at room temperature. An analysis by thin layer chromatography (TLC) indicated the presence of

starting material, and additional sodium borohydride (0.10 g, 2.6 mmol) was added. The reaction was stirred at room temperature overnight. The reaction was still incomplete, and sodium borohydride was added in portions (0.10 g and 0.20 g) until the starting material was consumed. The reaction mixture was filtered through a layer of CELITE filter agent.  
5 The filter cake was washed with dichloromethane until the filtrate was colorless, and the filtrate was then concentrated under reduced pressure. The crude product was purified by flash chromatography on silica gel (eluting sequentially with 30% ethyl acetate in hexane and 50% ethyl acetate in hexane) to provide 0.40 g of  $N^2,N^2$ -bis(4-methoxybenzyl)-6-pentyl- $N^4$ -(2-phenylethyl)pyridine-2,3,4-triamine as a yellow oil.

10 Part D

Carbonyl diimidazole (0.18 g, 1.1 mmol) was added to a solution of  $N^2,N^2$ -bis(4-methoxybenzyl)-6-pentyl- $N^4$ -(2-phenylethyl)pyridine-2,3,4-triamine (0.40 g, 0.74 mmol) in THF (4 mL), and the orange solution was heated at reflux for one hour and allowed to cool to room temperature. The volatiles were removed under reduced pressure, and the  
15 residue was purified by flash chromatography on silica gel (eluting sequentially with 30% ethyl acetate in hexane and 50% ethyl acetate in hexane) to provide 0.36 g of 4-[bis(4-methoxybenzyl)amino]-6-pentyl-1-(2-phenylethyl)-1*H*-imidazo[4,5-*c*]pyridin-2-ol as an oil that solidified upon standing.

Part E


20 A solution of 4-[bis(4-methoxybenzyl)amino]-6-pentyl-1-(2-phenylethyl)-1*H*-imidazo[4,5-*c*]pyridin-2-ol (0.36 g, 0.64 mmol) in trifluoroacetic acid (1.6 mL) was stirred at room temperature for five hours and then diluted with water (20 mL). The resulting mixture was adjusted to approximately pH 13 with the addition of aqueous sodium hydroxide (50% w/w) and stirred for one hour. A solid was present and was collected by  
25 filtration, washed with water, and recrystallized from acetonitrile (20 mL) and ethanol (5 mL). The crystals were washed with acetonitrile and dried under vacuum for 18 hours at 65 °C to provide 0.10 g of 4-amino-6-pentyl-1-(2-phenylethyl)-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a white crystalline powder, mp 238-240 °C.

$^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ )  $\delta$  10.10 (bs, 1H), 7.28-7.17 (m, 5H), 6.24 (s, 1H), 5.50 (bs, 2H), 3.92 (t,  $J = 7.5$  Hz, 2H) 2.90 (t,  $J = 7.5$  Hz, 2H), 2.44 (t,  $J = 7.5$  Hz, 2H), 1.55 (pentet,  $J = 7.5$  Hz, 2H), 1.31-1.21 (m, 4H), 0.87 (t,  $J = 7.5$  Hz, 3H);  
30 MS (APCI)  $m/z$  325 ( $M + H$ ) $^+$ ;

Anal. Calcd for  $C_{19}H_{24}N_4O$ : C, 70.34; H, 7.46; N, 17.27. Found: C, 70.21; H, 7.50; N, 17.31.

### Example 3

5 4-Amino-1-(2-hydroxy-2-methylpropyl)-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol



#### Part A

1-Amino-2-methylpropan-2-ol (0.64 g, 7.2 mmol) was added to a stirred solution of 2,4-dichloro-3-nitro-6-pentylpyridine (2.1 g, 8.0 mmol) and triethylamine (1.7 mL, 12 mmol) in DMF (40 mL). The reaction was stirred for 18 hours at room temperature and partitioned between water (200 mL) and ethyl acetate (50 mL). The organic layer was separated and washed with brine (50 mL), dried over sodium sulfate, filtered, and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel (eluting with 30% ethyl acetate in hexane) to provide 1.1 g of 1-[(2-chloro-3-nitro-6-pentylpyridin-4-yl)amino]-2-methylpropan-2-ol as a bright yellow oil.

#### Part B

1-[(2-Chloro-3-nitro-6-pentylpyridin-4-yl)amino]-2-methylpropan-2-ol (1.1 g, 3.5 mmol), triethylamine (0.7 mL, 5 mmol), and *N,N*-bis(4-methoxybenzyl)amine (1.3 g, 5.2 mmol) were reacted according to the method of Part B of Example 2 with the modification that the reaction was heated at reflux for six hours in toluene (35 mL) and then stirred at room temperature overnight to provide 1-({2-[bis(4-methoxybenzyl)amino]-3-nitro-6-pentylpyridin-4-yl}amino)-2-methylpropan-2-ol after the work-up procedure.

#### Part C

25 Sodium borohydride (0.100 g, 2.64 mmol) was added to a mixture of nickel(II) chloride hexahydrate (0.41 g, 1.7 mmol) in methanol (30 mL), and the mixture was stirred for 15 minutes. A solution of the material from Part B in methanol (28 mL) and dichloromethane (12 mL) was added. Additional sodium borohydride (0.140 g, 3.70 mmol) was added in portions, and the mixture was stirred for one hour at room



temperature. An analysis by TLC indicated the presence of starting material, and additional sodium borohydride was added in portions (0.12 g and 0.12 g). The reaction mixture was stirred briefly and then filtered through a layer of CELITE filter agent. The filter cake was washed with dichloromethane until the filtrate was colorless, and the filtrate was then concentrated under reduced pressure. The crude product was stirred with dichloromethane and filtered again through CELITE filter agent. The filtrate was concentrated under reduced pressure to provide 1-((3-amino-2-[bis(4-methoxybenzyl)amino]-6-pentylpyridin-4-yl)amino)-2-methylpropan-2-ol as a green oil.

#### Part D

The material from Part C was treated with carbonyl diimidazole (0.85 g, 5.2 mmol) according to the method described in Part D of Example 2 with the modification that chromatographic purification was carried out eluting sequentially with 50% ethyl acetate in hexane and then ethyl acetate. 4-[Bis(4-methoxybenzyl)amino]-1-(2-hydroxy-2-methylpropyl)-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol (1.4 g) was obtained as a colorless oil.

#### Part E

A solution of 4-[bis(4-methoxybenzyl)amino]-1-(2-hydroxy-2-methylpropyl)-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol (1.4 g, 2.6 mmol) in trifluoroacetic acid (7 mL) was stirred at room temperature for three hours and then diluted with water (20 mL). The resulting mixture was adjusted to approximately pH 13 with the addition of aqueous sodium hydroxide (50% w/w) and stirred for one hour. A solid was present and was collected by filtration and washed with water to provide 0.8 g of a white solid. The filtrate was allowed to stand for three days, and additional solid formed. The second solid was isolated by filtration and washed with water. The first solid was purified by column chromatography on silica gel (eluting with 10% methanol in dichloromethane) and then combined with the second solid. The combined solids were recrystallized from acetonitrile. The crystals were washed with acetonitrile and dried under vacuum for 17 hours at 65 °C to provide 0.35 g of 4-amino-1-(2-hydroxy-2-methylpropyl)-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol as white, crystalline plates, mp 240-243 °C.

<sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 10.16 (bs, 1H), 6.48 (s, 1H), 5.53 (bs, 2H), 4.58 (s, 1H), 3.59 (bs, 2H), 2.47 (t, *J* = 7.5 Hz, 2H), 1.59 (pentet, *J* = 7.5 Hz, 2H), 1.31-1.24 (m, 4H), 1.11 (bs, 6H), 0.85 (t, *J* = 7.5 Hz, 3H);


MS (APCI)  $m/z$  293 ( $M + H$ )<sup>+</sup>;

Anal. Calcd for C<sub>15</sub>H<sub>24</sub>N<sub>4</sub>O<sub>2</sub>: C, 61.62; H, 8.27; N, 19.16. Found: C, 61.49; H, 8.57; N, 19.25.

5

#### Example 4

4-Amino-6-pentyl-1-(tetrahydro-2H-pyran-4-ylmethyl)-1H-imidazo[4,5-c]pyridin-2-ol



#### Part A

10 Solid tetrahydro-2H-pyran-4-ylmethylamine hydrochloride (see U. S. Patent Application Publication No. 2004/0147543 (Hays *et al.*) Examples 477-480, 0.78 g, 5.1 mmol) was added to a stirred solution of 2,4-dichloro-3-nitro-6-pentylpyridine (1.5 g, 5.7 mmol) and triethylamine (2.0 mL, 14 mmol) in DMF (29 mL). The reaction was stirred for 18 hours at room temperature. The work-up and purification procedures described in Part A of Example 3 were followed to provide 1.0 g of 2-chloro-3-nitro-6-pentyl-*N*-  
15 (tetrahydro-2H-pyran-4-ylmethyl)pyridin-4-amine.

#### Part B

2-Chloro-3-nitro-6-pentyl-*N*-(tetrahydro-2H-pyran-4-ylmethyl)pyridin-4-amine (1.0 g, 2.9 mmol), triethylamine (0.61 mL, 4.4 mmol), and *N,N*-bis(4-methoxybenzyl)amine (1.1 g, 4.4 mmol) were reacted according to the method of Part B of  
20 Example 2 with the modification that the reaction was heated at reflux for 14 hours in toluene (29 mL) to provide *N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-3-nitro-6-pentyl-*N*<sup>4</sup>-(tetrahydro-2H-pyran-4-ylmethyl)pyridine-2,4-diamine as a yellow oil after the work-up procedure.

#### Part C

25 The method described in Part C of Example 3 was used to reduce the material from Part B to *N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-6-pentyl-*N*<sup>4</sup>-(tetrahydro-2H-pyran-4-ylmethyl)pyridine-2,3,4-triamine, which was obtained as a dark green oil.

#### Part D

The material from Part C was treated with carbonyl diimidazole (0.71 g, 4.4 mmol) according to the method described in Part D of Example 2 with the modification that

chromatographic purification was carried out using an automated flash chromatography system with a 40+M silica cartridge and eluting with 40% to 80% ethyl acetate in hexane. 4-[Bis(4-methoxybenzyl)amino]-6-pentyl-1-(tetrahydro-2H-pyran-4-ylmethyl)-1H-imidazo[4,5-c]pyridin-2-ol (0.75 g) was obtained as an oil that solidified upon standing.

5 Part E


A solution of 4-[bis(4-methoxybenzyl)amino]-6-pentyl-1-(tetrahydro-2H-pyran-4-ylmethyl)-1H-imidazo[4,5-c]pyridin-2-ol (0.75 g, 1.3 mmol) in trifluoroacetic acid (3 mL) was stirred at room temperature for 18 hours and then diluted with water (20 mL). The resulting mixture was adjusted to approximately pH 13 with the addition of aqueous sodium hydroxide (50% w/w) and stirred for one hour. A solid was present and was collected by filtration and washed with water to provide 0.5 g of a white solid. The filtrate was allowed to stand for three days, and additional solid formed. The second solid was isolated by filtration. The first solid was purified by automated flash chromatography (25+M silica cartridge, eluting with 0% to 15% methanol in dichloromethane) and then combined with the second solid. The combined solids (0.27 g) were recrystallized from acetonitrile (50 mL) and ethanol (8 mL). The crystals were washed with acetonitrile and dried under vacuum for 17 hours at 65 °C to provide 0.24 g of 4-amino-6-pentyl-1-(tetrahydro-2H-pyran-4-ylmethyl)-1H-imidazo[4,5-c]pyridin-2-ol as white, crystalline plates, mp >250 °C.

20 <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 10.13 (bs, 1H), 6.43 (s, 1H), 5.54 (bs, 2H), 3.83-3.79 (m, 2H), 3.58 (d, *J* = 7.5 Hz, 2H), 3.25-3.18 (m, 2H), 2.50-2.46 (m, 2H), 1.99-1.94 (m, 1H), 1.60 (pentet, *J* = 7.5 Hz, 2H), 1.47-1.43 (m, 2H), 1.32-1.22 (m, 6H), 0.86 (t, *J* = 6.9 Hz, 3H);

MS (APCI) *m/z* 319 (M + H)<sup>+</sup>;

25 Anal. Calcd for C<sub>17</sub>H<sub>26</sub>N<sub>4</sub>O<sub>2</sub>; C, 64.13; H, 8.23; N, 17.60. Found: C, 64.08; H, 8.11; N, 17.64.

## Example 5

4-Amino-6-pentyl-1-(pyridin-3-ylmethyl)-1*H*-imidazo[4,5-*c*]pyridin-2-ol

## Part A

5           3-(Aminomethyl)pyridine (0.52 mL, 5.1 mmol) was added to a stirred solution of 2,4-dichloro-3-nitro-6-pentylpyridine (1.5 g, 5.7 mmol) and triethylamine (2.0 mL, 14 mmol) in DMF (29 mL), and the methods described in Part A of Example 3 were followed with the modification that the crude product was purified by automated flash chromatography (40+M cartridge, eluting with 0% to 30% ethyl acetate in hexane) to provide 1.0 g of 2-chloro-3-nitro-6-pentyl-*N*-(pyridin-3-ylmethyl)pyridin-4-amine as a yellow oil.

## Part B

15           Triethylamine (0.63 mL, 4.5 mmol) and *N,N*-bis(4-methoxybenzyl)amine (1.2 g, 4.5 mmol) were added to a solution of 2-chloro-3-nitro-6-pentyl-*N*-(pyridin-3-ylmethyl)pyridin-4-amine (1.0 g, 3.0 mmol) in toluene (30 mL), and the yellow solution was heated at reflux for 14 hours, allowed to cool to room temperature, and concentrated under reduced pressure to provide *N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-3-nitro-6-pentyl-*N*<sup>4</sup>-(pyridin-3-ylmethyl)pyridine-2,4-diamine.

## Part C

20           Sodium borohydride (0.20 g, 5.4 mmol) was added to a mixture of nickel(II) chloride hexahydrate (0.36 g, 1.5 mmol) in methanol (20 mL), and the mixture was stirred for 15 minutes. A solution of the material from Part B in methanol (50 mL) and dichloromethane (11 mL) was added, and the reaction was stirred for 30 minutes. Additional sodium borohydride (0.20 g, 5.4 mmol) was added, and an analysis by TLC indicated the presence of starting material. Additional sodium borohydride was added (0.10 g), and then the reaction was complete. The reaction mixture was stirred briefly and then filtered through a layer of CELITE filter agent. The filter cake was washed with dichloromethane until the filtrate was colorless, and the filtrate was then concentrated under reduced pressure. The crude product was stirred with dichloromethane and filtered

again through CELITE filter agent. The filtrate was concentrated under reduced pressure to provide  $N^2,N^2$ -bis(4-methoxybenzyl)-6-pentyl- $N^4$ -(pyridin-3-ylmethyl)pyridine-2,3,4-triamine as a green oil.

#### Part D

5           The material from Part C was treated with carbonyl diimidazole (0.73 g, 4.5 mmol) according to the method described in Part D of Example 2 with the modification that chromatographic purification was carried out using an automated flash chromatography system with a 40+M silica cartridge and eluting with a gradient of 70% ethyl acetate in hexane to 100% ethyl acetate. 4-[Bis(4-methoxybenzyl)amino]-6-pentyl-1-(pyridin-3-ylmethyl)-1*H*-imidazo[4,5-*c*]pyridin-2-ol (1.3 g) was obtained as an oil that solidified  
10           somewhat upon standing.

#### Part E


          A solution of 4-[bis(4-methoxybenzyl)amino]-6-pentyl-1-(pyridin-3-ylmethyl)-1*H*-imidazo[4,5-*c*]pyridin-2-ol (1.3 g, 2.4 mmol) in trifluoroacetic acid (6 mL) was stirred at  
15           room temperature for two hours and then diluted with water (20 mL). The resulting mixture was adjusted to approximately pH 13 with the addition of aqueous sodium hydroxide (50% w/w) and stirred for two hours. A solid was present and was collected by filtration and washed with water to provide 1 g of a white solid. The solid was purified by automated flash chromatography (40+M silica cartridge, eluting with 0% to 20% methanol  
20           in dichloromethane) followed by recrystallization from ethanol (10 mL). The crystals were washed with ethanol and dried under vacuum for four hours at 65 °C to provide 4-amino-6-pentyl-1-(pyridin-3-ylmethyl)-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a white crystalline powder, mp 243-245 °C.

$^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ )  $\delta$  10.28 (bs, 1H), 8.57 (d,  $J = 2.5$  Hz, 1H), 8.48-8.47 (m, 1H), 7.68-7.66 (m, 1H), 7.37-7.34 (m, 1H), 6.41 (s, 1H), 5.59 (bs, 2H), 4.96 (bs, 2H), 2.45  
25           (t,  $J = 7.5$  Hz, 2H), 1.56 (pentet,  $J = 7.5$  Hz, 2H), 1.28-1.19 (m, 4H), 0.83 (t,  $J = 6.9$  Hz, 3H);

          MS (APCI)  $m/z$  312 ( $M + H$ ) $^+$ ;


          Anal. Calcd for  $\text{C}_{17}\text{H}_{21}\text{N}_5\text{O} \cdot 0.25 \text{H}_2\text{O}$ : C, 64.64; H, 6.86; N, 22.17. Found: C, 64.32; H,  
30           7.14; N, 22.16.

## Example 6

4-Amino-1-benzyl-6-methyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol

4-Amino-1-benzyl-6-methyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol was prepared according to the general methods described in Parts G through K of Example 1 using 2,4-dichloro-6-methyl-3-nitropyridine in lieu of 2,4-dichloro-3-nitro-6-pentylpyridine in Part G. The crude product was purified by automated flash chromatography (40+M cartridge, eluting with 0% to 15% methanol in dichloromethane) followed by recrystallization from methanol to provide 4-amino-1-benzyl-6-methyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a crystalline solid, mp > 300 °C. Anal. Calcd for C<sub>14</sub>H<sub>14</sub>N<sub>4</sub>O: C, 66.13; H, 5.55; N, 22.03. Found: C, 66.22; H, 5.33; N, 22.03.

## Example 7

4-Amino-6-pentyl-1-{2-[3-[(1,3-thiazolo-2-yl)propoxy]ethyl]-1*H*-imidazo[4,5-*c*]pyridin-2-ol


4-Amino-6-pentyl-1-{2-[3-[(1,3-thiazolo-2-yl)propoxy]ethyl]-1*H*-imidazo[4,5-*c*]pyridin-2-ol was prepared according to the general methods described in Parts A through E of Example 2 using 2-[3-[(1,3-thiazolo-2-yl)propoxy]ethylamine (see U.S. Patent No. 6,797,718 (Dellaria *et al.*) Example 82) in lieu of phenethylamine in Part A. The crude product was purified by automated flash chromatography (40+M cartridge, eluting with 0% to 20% methanol in dichloromethane) followed by recrystallization from acetonitrile

to provide 4-amino-6-pentyl-1-{2-[3-[(1,3-thiazolo-2-yl)propoxy]ethyl]-1*H*-imidazo[4,5-*c*]pyridin-2-yl} as a crystalline solid, mp 177.0-178.0 °C. Anal. Calcd for C<sub>19</sub>H<sub>27</sub>N<sub>5</sub>O<sub>2</sub>S: C, 58.59; H, 6.99; N, 17.98. Found: C, 58.66; H, 6.86; N, 18.03.

5

## Example 8

*N*-[3-(4-Amino-2-hydroxy-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-1-yl)propyl]methanesulfonamide



## Part A

10        *Tert*-butyl 3-aminopropylcarbamate (2.4 g, 13.68 mmol) was added to a solution of 2,4-dichloro-3-nitro-6-pentylpyridine (4 g, 15.2 mmol) and triethylamine (3.8 g, 38.0 mmol) in DMF (76 mL). The reaction mixture was stirred at ambient temperature for 18 hours and then partitioned between ethyl acetate (50 mL) and water (200 mL). The organic layer was washed with brine (50 mL), dried over sodium sulfate, filtered, and then concentrated under reduced pressure to provide crude product as a bright yellow oil. This material was purified by automated flash chromatography (40+M cartridge, eluting with 10% to 50% ethyl acetate in hexanes) to provide 3 g of *tert*-butyl 3-[(2-chloro-3-nitro-6-pentylpyridin-4-yl)amino]propylcarbamate as a yellow oil.

## Part B

20        A solution of the material from Part A (3 g, 7.48 mmol), *N,N*-bis(4-methoxybenzyl)amine (2.9 g, 11.22 mmol), and triethylamine (1.1 g, 11.22 mmol) in toluene (75 mL) was heated at reflux for 18 hours and then concentrated under reduced pressure to provide 4.6 g of crude *tert*-butyl 3-[(2-[bis(4-methoxybenzyl)amino]-3-nitro-6-pentylpyridin-4-yl)amino]propylcarbamate.

## 25        Part C

Solid sodium borohydride (0.25 g, 6.6 mmol) was added in a single portion to a solution of nickel(II) chloride hexahydrate (0.9 g, 3.7 mmol) in methanol (50 mL) and the

resulting suspension was stirred for 15 minutes. A solution of the material from Part B (about 7.5 mmol) in a mixture of dichloromethane (27 mL) and methanol (75 mL) was added in a single portion to the suspension. Sodium borohydride (0.26 g, 6.9 mmol) was added. After 30 minutes more sodium borohydride (0.2 g, 5 mmol) was added. Small portions of sodium borohydride were added until analysis by thin layer chromatography indicated that all of the starting material had been consumed. The reaction mixture was filtered through a layer of CELITE filter agent and the filter cake was washed with dichloromethane until the wash was clear. The filtrate was concentrated under reduced pressure to provide about 4.4 g of *tert*-butyl 3-({3-amino-2-[bis(4-methoxybenzyl)amino]-6-pentylpyridin-4-yl}amino)propylcarbamate as a green oil.

#### Part D

A solution of the material from Part C (about 7.5 mmol) and carbonyldiimidazole (1.8 g, 11 mmol) in THF (37 mL) was heated at reflux for 1 hour, cooled to ambient temperature, and then concentrated under reduced pressure to provide crude product. This material was purified by automated flash chromatography (40+M cartridge, eluting with 70% to 100% ethyl acetate in hexanes) to provide 3 g of *tert*-butyl 3-{4-[bis(4-methoxybenzyl)amino]-2-hydroxy-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-1-yl}propylcarbamate as a solid.

#### Part E

A solution of the material from Part D in trifluoroacetic acid (12 mL) was stirred at ambient temperature for 2 hours. Water (20 mL) was added and a white precipitate formed. The pH was adjusted to about 13 with 50% sodium hydroxide and the suspension was stirred at ambient temperature for 96 hours. The solid was removed by filtration and the filter cake was rinsed with water. The filtrate was concentrated under reduced pressure to provide a solid. This material was slurried with chloroform (100 mL) for 1 hour. The solid was removed by filtration. The filtrate was concentrated under reduced pressure to provide crude product as a brown foam. The foam was purified by automated flash chromatography (40+M cartridge, eluting with 30% to 60% CMA in chloroform) to provide 0.6 g of 4-amino-1-(3-aminopropyl)-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a white, waxy solid.




## Part F

Methanesulfonyl chloride (0.12 g, 1.08 mmol) was added to a suspension of material from Part E (0.25 g, 0.90 mmol) in triethylamine (0.31 mL) and dichloromethane (5 mL). The resulting solution was stirred at ambient temperature for 1 hour and then partitioned between dichloromethane (50 mL) and saturated aqueous ammonium chloride (50 mL). The organic layer was dried over sodium sulfate, filtered, and then concentrated under reduced pressure to provide crude product as a solid. This material was purified by automated flash chromatography (25+M cartridge, eluting with 0% to 10% methanol in dichloromethane) followed by recrystallization from a mixture of acetonitrile (15 mL) and ethanol (2 mL) to provide 0.2 g of *N*-[3-(4-amino-2-hydroxy-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-1-yl)propyl]methanesulfonamide as crystalline solid, mp 174.0-176.0 °C. Anal. Calcd for C<sub>15</sub>H<sub>25</sub>N<sub>5</sub>O<sub>3</sub>S: C, 50.68; H, 7.09; N, 19.70. Found: C, 50.76; H, 7.42; N, 19.85.

## Example 9


*N*-[3-(4-Amino-2-hydroxy-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-1-yl)propyl]-*N'*-cyclohexylurea



Cyclohexyl isocyanate (0.1 g, 0.8 mmol) was added to a suspension of 4-amino-1-(3-aminopropyl)-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol (0.2 g, 0.7 mmol) in dichloromethane (4 mL). The resulting solution was stirred at ambient temperature for 2 hours and then concentrated under reduced pressure to provide crude product. This material was purified by automated flash chromatography (25+M cartridge, eluting with 0% to 10% methanol in dichloromethane) followed by recrystallization from a mixture of acetonitrile (15 mL) and ethanol (8 mL) to provide 0.2 g of *N*-[3-(4-amino-2-hydroxy-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-1-yl)propyl]-*N'*-cyclohexylurea as a crystalline solid, mp 202.0-204.0 °C. Anal. Calcd for C<sub>21</sub>H<sub>34</sub>N<sub>6</sub>O<sub>2</sub>: C, 62.66; H, 8.51; N, 20.88. Found: C, 62.55; H, 8.88; N, 20.86.

## Example 10


*N*-[3-(4-Amino-2-hydroxy-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-1-yl)propyl]acetamide



Acetyl chloride (0.05 g, 0.60 mmol) was added dropwise to a suspension of 4-amino-1-(3-aminopropyl)-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol (0.15 g, 0.54 mmol) in triethylamine (0.19 mL, 1.5 mmol) and dichloromethane (3 mL). The reaction mixture was stirred at ambient temperature for 2 hours and then partitioned between dichloromethane (50 mL) and saturated aqueous ammonium chloride (50 mL). The organic layer was dried over sodium sulfate, filtered, and then concentrated under reduced pressure to provide crude product as an oil. This material was purified by automated flash chromatography (25+M cartridge, eluting with 0% to 15% methanol in dichloromethane) followed by recrystallization from a mixture of acetonitrile (15 mL) and ethanol (3 mL) to provide 0.1 g of *N*-[3-(4-amino-2-hydroxy-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-1-yl)propyl]acetamide as a crystalline solid, mp 189.0-191.0 °C. Anal. Calcd for C<sub>16</sub>H<sub>25</sub>N<sub>5</sub>O<sub>2</sub>: C, 60.17; H, 7.89; N, 21.93. Found: C, 59.96; H, 7.95; N, 21.98.

## Example 11

4-Amino-6-pentyl-1-(piperidin-4-ylmethyl)-1*H*-imidazo[4,5-*c*]pyridin-2-ol



## 20 Part A

A solution of 4-(aminomethyl)piperidine (10 g, 87.6 mmol) in chloroform (60 mL) was chilled in an ice water bath. A solution of di-*tert*-butyl dicarbonate (9.6 g, 43.8 mmol) in chloroform (37 mL) was added dropwise over a period of 30 minutes. The ice bath was removed and the reaction mixture was stirred at ambient temperature over the weekend. The reaction was quenched with water (100 mL). The organic layer was dried

over sodium sulfate, filtered, and then concentrated under reduced pressure to provide 9 g of *tert*-butyl 4-(aminomethyl)piperidine-1-carboxylate.

#### Part B

*Tert*-butyl 4-(aminomethyl)piperidine-1-carboxylate (2.2 g, 10.3 mmol) was added dropwise to a solution of 2,4-dichloro-3-nitro-6-pentylpyridine (3 g, 11.4 mmol) and triethylamine (1.7 g, 17.1 mmol) in DMF (57 mL). The reaction mixture was stirred at ambient temperature for 18 hours then quenched with water (200 ml) and extracted with ethyl acetate (2 x 50 mL). The organics were combined, washed with brine (50 mL), dried over sodium sulfate, filtered, and then concentrated under reduced pressure to provide crude product as an oil. This material was purified by automated flash chromatography (40+M cartridge, eluting with 0% to 10% ethyl acetate in hexanes) to provide 2.9 g of *tert*-butyl 4-[[2-chloro-3-nitro-6-pentylpyridin-4-yl]amino]methyl}piperidine-1-carboxylate.

#### Part C

A solution of the material from Part B (2.9 g, 6.58 mmol), *N,N*-bis(4-methoxybenzyl)amine (2.5 g, 9.87 mmol), and triethylamine (1 g, 9.87 mmol) in toluene (66 mL) was heated at reflux for 22 hours and then concentrated under reduced pressure to provide about 4.4 g of *tert*-butyl 4-[(2-[bis(4-methoxybenzyl)amino]-3-nitro-6-pentylpyridin-4-yl)amino)methyl]piperidine-1-carboxylate.

#### Part D

Solid sodium borohydride (0.2 g, 5 mmol) was added in a single portion to a solution of nickel(II) chloride hexahydrate (0.78 g, 3.3 mmol) in methanol (50 mL) and the resulting suspension was stirred for 15 minutes. A solution of the material from Part C (about 6.6 mmol) in a mixture of dichloromethane (24 mL) and methanol (60 mL) was added in a single portion to the suspension. Sodium borohydride (0.25 g, 6.6 mmol) was added. After 30 minutes more sodium borohydride (0.2 g, 5 mmol) was added. Small portions of sodium borohydride were added until analysis by thin layer chromatography indicated that all of the starting material had been consumed. The reaction mixture was filtered through a layer of CELITE filter agent and the filter cake was washed with dichloromethane until the wash was clear. The filtrate was concentrated under reduced pressure. The residue was triturated with dichloromethane then filtered through a layer of CELITE filter agent. The filtrate was concentrated under reduced pressure to provide

about 4.2 g of *tert*-butyl 4-[(3-amino-2-[bis(4-methoxybenzyl)amino]-6-pentylpyridin-4-yl)methyl]piperidine-1-carboxylate as a green oil.

#### Part E


A solution of the material from Part D (about 6.6 mmol) and carbonyldiimidazole (1.6 g, 9.9 mmol) in THF (33 mL) was heated at reflux for 1 hour, cooled to ambient temperature, and then concentrated under reduced pressure to provide crude product. This material was purified by automated flash chromatography (40+M cartridge, eluting with 60% to 100% ethyl acetate in hexanes) to provide 4 g of *tert*-butyl 4-[(4-[bis(4-methoxybenzyl)amino]-2-hydroxy-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-1-yl)methyl]piperidine-1-carboxylate as a solid.

#### Part F

A solution of the material from Part E in trifluoroacetic acid (15 mL) was stirred at ambient temperature for 18 hours. Water (20 mL) was added and the pH was adjusted to about 14 with 50% sodium hydroxide. The mixture was neutralized (pH 7) with 1M hydrochloric acid. The resulting suspension was stirred for 1 hour. The solid was isolated by filtration, washed with water, and dried. This material was purified by automated flash chromatography (40+M cartridge, eluting with 50% to 100% CMA in chloroform) followed by recrystallization from ethanol (22 mL) to provide 1.2 g of 4-amino-6-pentyl-1-(piperidin-4-ylmethyl)-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a crystalline solid, mp 246.0-249.0 °C. Anal. Calcd for C<sub>17</sub>H<sub>27</sub>N<sub>5</sub>O: C, 64.32; H, 8.57; N, 22.06. Found: C, 64.10; H, 8.62; N, 21.87.

#### Example 12

4-Amino-1-{[1-(methylsulfonyl)piperidin-4-yl]methyl}-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol




25

Solid methanesulfonic anhydride (0.31 g, 1.78 mmol) was added in a single portion to a suspension of 4-amino-6-pentyl-1-(piperidin-4-ylmethyl)-1*H*-imidazo[4,5-*c*]pyridin-2-ol (0.47 g, 1.48 mmol) in dichloromethane (6 mL). The reaction mixture was stirred at

ambient temperature for 2 hours. More methanesulfonic anhydride (0.05 g) was added. The reaction mixture was stirred for an additional 30 minutes and then quenched with saturated aqueous sodium carbonate (20 mL) and allowed to stir overnight. A solid was isolated by filtration, washed with water, and then recrystallized from ethanol to provide 0.4 g of 4-amino-1-[[1-(methanesulfonyl)piperidin-4-yl]methyl]-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a solid, mp 228.0-231.0 °C. Anal. Calcd for C<sub>18</sub>H<sub>29</sub>N<sub>5</sub>O<sub>3</sub>S: C, 54.66; H, 7.39; N, 17.71. Found: C, 54.45; H, 7.38; N, 17.65.


## Example 13

1-[[1-(1-Acetylpiperidin-4-yl)methyl]-4-amino-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol




Acetyl chloride (0.15 g, 1.91 mmol) was added to a suspension of 4-amino-6-pentyl-1-(piperidin-4-ylmethyl)-1*H*-imidazo[4,5-*c*]pyridin-2-ol (0.55 g, 1.73 mmol) and triethylamine (0.44 g, 4.33 mmol) in dichloromethane (9 mL). The reaction mixture was stirred at ambient temperature for 2 hours. The reaction mixture was loaded directly onto a silica cartridge and purified by automated flash chromatography (25+M cartridge, eluting with 0% to 15% methanol in dichloromethane) to provide product as a white solid. This material was recrystallized from acetonitrile to provide 0.23 g of 1-[[1-(1-acetylpiperidin-4-yl)methyl]-4-amino-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a crystalline solid, mp 228.0-231.0 °C. Anal. Calcd for C<sub>19</sub>H<sub>29</sub>N<sub>5</sub>O<sub>2</sub>: C, 63.48; H, 8.13; N, 19.48. Found: C, 63.22; H, 8.21; N, 19.33.

## Example 14

4-Amino-1-{[1-(ethylsulfonyl)piperidin-4-yl]methyl}-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol

5           Ethanesulfonyl chloride (0.15 g, 1.13 mmol) was added to a suspension of 4-amino-6-pentyl-1-(piperidin-4-ylmethyl)-1*H*-imidazo[4,5-*c*]pyridin-2-ol (0.3 g, 0.94 mmol) and triethylamine (0.24 g, 2.36 mmol) in dichloromethane (5 mL). The reaction mixture was stirred at ambient temperature for 2 hours. The reaction mixture was quenched with methanol, which brought all of the solids into solution. The solution was  
10           loaded directly onto a silica cartridge and purified by automated flash chromatography (25+M cartridge, eluting with 0% to 10% methanol in dichloromethane) to provide 0.13 g of product as a white solid. This material was recrystallized from ethanol (7 mL) to provide 0.1 g 4-amino-1-{[1-(ethylsulfonyl)piperidin-4-yl]methyl}-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a crystalline solid, mp 222.0-225.0 °C. Anal. Calcd for  
15           C<sub>19</sub>H<sub>31</sub>N<sub>5</sub>O<sub>3</sub>S: C, 55.72; H, 7.63; N, 17.10. Found: C, 55.91; H, 7.53; N, 17.04.


## Example 15

4-Amino-1-{[1-(cyclopropylcarbonyl)piperidin-4-yl]methyl}-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol

20           Cyclopropanecarbonyl chloride (0.12 g, 1.13 mmol) was added dropwise to a suspension of 4-amino-6-pentyl-1-(piperidin-4-ylmethyl)-1*H*-imidazo[4,5-*c*]pyridin-2-ol (0.3 g, 0.94 mmol) and triethylamine (0.24 g, 2.36 mmol) in dichloromethane (5 mL). The reaction mixture was stirred at ambient temperature for 1 hour. The reaction mixture was

quenched with methanol, which brought all of the solids into solution. The solution was loaded directly onto a silica cartridge and purified by automated flash chromatography (25+M cartridge, eluting with 0% to 10% methanol in dichloromethane) to provide 0.13 g of product as a white solid. This material was recrystallized from ethanol (7 mL) to provide 0.2 g of 4-amino-1-[[1-(cyclopropylcarbonyl)piperidin-4-yl]methyl]-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a crystalline solid, mp 205.0-207.0 °C. Anal. Calcd for C<sub>21</sub>H<sub>31</sub>N<sub>5</sub>O<sub>2</sub>: C, 65.43; H, 8.11; N, 18.17. Found: C, 65.30; H, 8.03; N, 18.22.

## Example 16

4-Amino-1-[4-(methylsulfonyl)butyl]-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol

## Part A

4-(Methylsulfonyl)butan-1-amine (1.9 g of 50% pure material) was added to a solution of 2,4-dichloro-3-nitro-6-pentylpyridine (1.5 g, 5.70 mmol) and triethylamine (1.4 g, 14.2 mmol) in DMF (29 mL). The reaction mixture was stirred at ambient temperature for 18 hours and then partitioned between ethyl acetate (50 mL) and water (200 mL). The organic layer was washed with brine (50 mL), dried over sodium sulfate, filtered, and then concentrated under reduced pressure to provide crude product as a bright yellow oil. This material was purified by automated flash chromatography (40+M cartridge, eluting with 30% to 80% ethyl acetate in hexanes) to provide 1.3 g of 2-chloro-*N*-[4-(methylsulfonyl)butyl]-3-nitro-6-pentylpyridin-4-amine as a yellow oil.

## Part B

A solution of the material from Part A (1.3 g, 3.44 mmol), *N,N*-bis(4-methoxybenzyl)amine (1.3 g, 5.16 mmol), and triethylamine (0.5 g, 5.16 mmol) in toluene (34 mL) was heated at reflux for 22 hours and then concentrated under reduced pressure to provide about 2.1 g of *N,N*-bis(4-methoxybenzyl)-*N*^4-[4-(methylsulfonyl)butyl]-3-nitro-6-pentylpyridine-2,4-diamine.

## Part C

Solid sodium borohydride (0.1 g, 2.6 mmol) was added in a single portion to a solution of nickel(II) chloride hexahydrate (0.41 g, 1.7 mmol) in methanol (30 mL) and the resulting suspension was stirred for 15 minutes. A solution of the material from Part B (about 3.4 mmol) in a mixture of dichloromethane (12 mL) and methanol (27 mL) was added in a single portion to the suspension. Sodium borohydride (0.13 g, 3.3 mmol) was added. After 30 minutes more sodium borohydride (0.2 g, 5 mmol) was added. Small portions of sodium borohydride were added until analysis by thin layer chromatography indicated that all of the starting material had been consumed. The reaction mixture was filtered through a layer of CELITE filter agent and the filter cake was washed with dichloromethane until the wash was clear. The filtrate was concentrated under reduced pressure. The residue was triturated with dichloromethane then filtered through a layer of CELITE filter agent. The filtrate was concentrated under reduced pressure to provide about 2 g of *N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-*N*<sup>4</sup>-[4-(methylsulfonyl)butyl]-6-pentylpyridine-2,3,4-triamine as a green oil.

## Part D

A solution of the material from Part C (about 3.4 mmol) and carbonyldiimidazole (0.84 g, 5.2 mmol) in THF (17 mL) was heated at reflux for 1 hour, cooled to ambient temperature, and then concentrated under reduced pressure to provide crude product. This material was purified by automated flash chromatography (40+M cartridge, eluting with 0% to 5% methanol in dichloromethane) to provide an oil. The oil was purified by automated flash chromatography (40+M cartridge, eluting with 70% to 100% ethyl acetate in hexanes) to provide 1.6 g of 4-[bis(4-methoxybenzyl)amino]-1-[4-(methylsulfonyl)butyl]-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a white solid.

## Part E

A solution of the material from Part D in trifluoroacetic acid (7 mL) was stirred at ambient temperature for 2 hours. Water (20 mL) was added and the pH was adjusted to about 12 with 50% sodium hydroxide. The resulting suspension was stirred for 1 hour. The solid was isolated by filtration and washed with water to provide about 1 g of crude product as a solid. This material was purified by automated flash chromatography (40+M cartridge, eluting with 0% to 10% methanol in dichloromethane) followed by recrystallization from acetonitrile to provide 0.56 g of 4-amino-1-[4-




(methylsulfonyl)butyl]-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a crystalline solid, mp 188.0-189.0 °C. Anal. Calcd for C<sub>16</sub>H<sub>26</sub>N<sub>4</sub>O<sub>3</sub>S: C, 54.21; H, 7.39; N, 15.81. Found: C, 54.08; H, 7.51; N, 15.77.

5

## Example 17

4-Amino-1-[2-(methylsulfonyl)ethyl]-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol




4-Amino-1-[2-(methylsulfonyl)ethyl]-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol was prepared according to the methods described in Parts A through E of Example 16 using 2-(methylsulfonyl)ethan-1-amine in lieu of 4-(methylsulfonyl)butan-1-amine in Part A. The crude product was purified by automated flash chromatography (40+M cartridge, eluting with 0% to 15% methanol in dichloromethane) followed by recrystallization from acetonitrile to provide 0.65 g of 4-amino-1-[2-(methylsulfonyl)ethyl]-6-pentyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a crystalline solid, mp 223.0-225.0 °C. Anal. Calcd for C<sub>14</sub>H<sub>22</sub>N<sub>4</sub>O<sub>3</sub>S: C, 51.51; H, 6.79; N, 17.16. Found: C, 51.80; H, 6.95; N, 17.20.

15

## Example 18

4-Amino-6-(2-ethoxyethyl)-1-[4-(methylsulfonyl)butyl]-1*H*-imidazo[4,5-*c*]pyridin-2-ol



20

## Part A

To a solution of 3-ethoxypropionic acid (35.0 g, 296 mmol) and 1-(methylsulfonyl)benzotriazole (58.3 g, 296 mmol) in THF (400 mL) was added triethylamine (57.7 mL, 414 mmol). The resultant solution was heated to reflux overnight

under a nitrogen atmosphere. The following morning, solvents were removed via rotary evaporation, and the residue was partitioned between  $\text{CH}_2\text{Cl}_2$  and aqueous 1N HCl. The aqueous phase was extracted with  $\text{CH}_2\text{Cl}_2$  (2x). The combined organic layers were washed with brine and dried over  $\text{MgSO}_4$ , then filtered through a silica gel plug, eluting with 1L of 2% MeOH /  $\text{CH}_2\text{Cl}_2$ . Removal of the solvents by rotary evaporation afforded about 65 g of 1-(3-ethoxypropanoyl)-1*H*-1,2,3-benzotriazole as a clear, pale yellow oil that solidified upon standing. Analysis by  $^1\text{H}$  NMR revealed product of sufficient purity to carry forward without additional purification.

#### Part B

A 1L round-bottomed flask was charged with sodium hydride (11.7 g of a 60% dispersion in oil, 293 mmol). The sodium hydride was washed with hexanes (2x); then THF (300 mL) was added to the flask. A solution of ethyl acetoacetate (34.6 g, 266 mmol) in THF (100 mL) was then added dropwise via addition funnel under a nitrogen atmosphere. A thick white precipitate formed at this point. After stirring for 1 hour, a solution of 1-(3-ethoxypropanoyl)-1*H*-1,2,3-benzotriazole (58.3 g, 266 mmol) in THF (100 mL) was added via addition funnel. The solution became homogeneous, then eventually a cloudy yellow mixture. This mixture was allowed to stir at room temperature under a nitrogen atmosphere overnight. The following morning, a solution of ammonium chloride (47.0 g, 878 mmol) and ammonium hydroxide (11.2 mL) in de-ionized water (50 mL) was added, and the resultant solution was heated to reflux for 2 hours. The volatile solvents were then removed by rotary evaporation, and the remaining aqueous layer was adjusted to pH 4-5 by addition of aqueous 1N HCl. The aqueous layer was then extracted with EtOAc (4 x 200 mL), and the combined organic layers were washed with brine, dried over  $\text{MgSO}_4$ , filtered, and concentrated to a yellow oil. This material was purified by suction filter chromatography on silica gel, eluting with 3/1 hexane/EtOAc, to afford 40.2 g (80% yield) of ethyl 5-ethoxy-3-oxopentanoate as a yellow oil. Analysis by  $^1\text{H}$  NMR revealed material of sufficient purity to carry forward.

#### Part C

To a solution of ethyl 5-ethoxy-3-oxopentanoate (40.2 g, 214 mmol) in methanol (80 mL) was added ammonium acetate (82.3 g, 1.07 mol). The resultant solution was allowed to stir at room temperature for 72 hours. The methanol was then removed by rotary evaporation, and chloroform was added to the residue. The white precipitate that

formed was removed via filtration through a fritted glass funnel, and the filtrate was washed with water (2x) and brine (1x), then dried over MgSO<sub>4</sub>, filtered, and concentrated to afford 41 g of ethyl 3-amino-5-ethoxypent-2-enoate as a yellow oil. Analysis of this material by <sup>1</sup>H NMR revealed it to be clean product, which was carried on without further purification.

#### Part D

A solution of ethyl 3-amino-5-ethoxypent-2-enoate (40.1 g, 214 mmol) and pyridine (20.3 g, 257 mmol) in THF (400 mL) was cooled in an ice bath, and a solution of methyl malonyl chloride (32.2 g, 236 mmol) in THF (100 mL) was added dropwise via addition funnel under a nitrogen atmosphere. Upon complete addition, the resultant mixture was warmed to room temperature and allowed to stir overnight under a nitrogen atmosphere. The following morning, the mixture was diluted with water and extracted with EtOAc (3 x 200 mL). The combined organic layers were washed with brine, dried over MgSO<sub>4</sub>, filtered, and concentrated to afford about 62 g of ethyl 5-ethoxy-3-[(3-methoxy-3-oxopropanoyl)amino]pent-2-enoate as a yellow oil. Analysis of this material by <sup>1</sup>H NMR revealed the desired product, along with unidentified impurities. The material was carried on without further purification.

#### Part E

A 1L round-bottomed flask was charged with sodium hydride (17.1 g of a 60% dispersion in oil, 428 mmol). The sodium hydride was washed with hexanes (2x); then THF (400 mL) was added to the flask. A solution of ethyl 5-ethoxy-3-[(3-methoxy-3-oxopropanoyl)amino]pent-2-enoate (61.5 g, 214 mmol) in THF (150 mL) was then added dropwise via addition funnel under a nitrogen atmosphere. Hydrogen evolution was apparent, after which time the mixture became a thick gel. Additional THF (100 mL) was added, and the reaction mixture was heated to reflux for 4 hours under a nitrogen atmosphere. The reaction was then cooled to room temperature and quenched by careful addition of methanol. The mixture was then diluted with water, and the pH was adjusted to about 4 by addition of 1N aqueous HCl. This mixture was extracted with dichloromethane (4 x 200 mL), and the combined organic layers were washed with brine, dried over MgSO<sub>4</sub>, filtered, and concentrated to afford a yellow solid. This material was triturated with 1/1 hexane/EtOAc to afford 23.4 g (43% yield) of ethyl 2-(2-ethoxyethyl)-

4-hydroxy-6-oxo-1,6-dihydropyridine-3-carboxylate as a light yellow solid. Purity was established by  $^1\text{H}$  NMR.

#### Part F

The material from Part E (23.4 g, 91.7 mmol) was dissolved in 70 mL of aqueous  
5 3N HCl, and the resultant solution was heated to reflux for 24 hours. Upon cooling to room temperature, the pH of the solution was adjusted to 7 by addition of ammonium hydroxide. The water was removed by rotary evaporation, and methanol was added to the residue. The mixture was filtered through a layer of CELITE filter agent, and the solvents were removed by rotary evaporation. The residue was adsorbed onto silica gel and placed  
10 on top of a short silica gel plug. The desired product was flushed through this column, ramping the eluent from 10-30% MeOH in  $\text{CH}_2\text{Cl}_2$ . This provided 2.70 g (16% yield) of 6-(2-ethoxyethyl)pyridine-2,4-diol as a pale yellow solid. Purity was established by  $^1\text{H}$  NMR.

#### Part G

The material from Part F (2.70 g, 14.7 mmol) was dissolved in acetic acid (10 mL),  
15 and nitric acid (1.4 mL, 22.1 mmol) was added via syringe. The resultant dark solution was heated in an 85 °C oil bath for 2 hours, during which time the color became a light yellow-green. After cooling to room temperature, toluene (15 mL) was added to the solution, and solvents were removed via rotary evaporation. The residue was dissolved in  
20 MeOH, and the pH was adjusted to 7-8 by addition of ammonium hydroxide. Silica gel was added to the solution, and solvents were removed by rotary evaporation. The silica gel containing the adsorbed product was loaded onto a silica gel column, and the product was eluted with a solvent system ramped from 3/1 to 1/1  $\text{CH}_2\text{Cl}_2/\text{MeOH}$ . This afforded 1.93 g (58% yield) of 6-(2-ethoxyethyl)-3-nitropyridine-2,4-diol as a yellow solid. Purity  
25 was established by LC-MS (229 = M+H) and  $^1\text{H}$  NMR.

#### Part H

6-(2-Ethoxyethyl)-3-nitropyridine-2,4-diol (2.42 g, 10.6 mmol) was dissolved in  
POCl<sub>3</sub> (36.0 mL, 386 mmol), and the resultant yellow solution was heated in an 80 °C oil  
bath. Over several hours, the solution slowly turned dark in color. The bulk of the POCl<sub>3</sub>  
30 was removed by rotary evaporation, and the residue was quenched by careful addition of water. The pH was adjusted to 9 by addition of Na<sub>2</sub>CO<sub>3</sub>, and the mixture was then extracted with  $\text{CH}_2\text{Cl}_2$  (3 x 50 mL). The combined organic layers were washed with brine,

dried over  $\text{MgSO}_4$ , filtered, and concentrated to a dark oil. Purification via flash chromatography on silica gel (4/1 hexane/EtOAc eluent) afforded 1.23 g (44% yield) of 2,4-dichloro-6-(2-ethoxyethyl)-3-nitropyridine as a slightly tan oil that was quite clean by  $^1\text{H}$  NMR analysis.

5     Part I

The material from Part H (1.23 g, 4.64 mmol) was dissolved in  $\text{CH}_2\text{Cl}_2$  (100 mL), and 4-(methylsulfonyl)butan-1-amine (1.54 g, 10.2 mmol) and triethylamine (1.62 mL, 11.6 mmol) were added. The resultant solution was allowed to stir at room temperature overnight under a nitrogen atmosphere. The following morning, the solution was washed  
10     with saturated aqueous  $\text{NaHCO}_3$  and brine, then dried over  $\text{MgSO}_4$ , filtered, and concentrated to a yellow oil. Purification via flash chromatography on silica gel (ramp eluent from 2-3% MeOH in  $\text{CH}_2\text{Cl}_2$ ) afforded 1.13 g (64% yield) of 2-chloro-6-(2-ethoxyethyl)-*N*-[4-(methylsulfonyl)butyl]-3-nitropyridin-4-amine as a thick yellow oil that was clean by  $^1\text{H}$  NMR analysis.

15     Part J

The material from Part I (1.13 g, 2.97 mmol) was dissolved in toluene (60 mL), and triethylamine (0.62 mL, 4.5 mmol) and di-*para*-methoxy benzylamine (1.15 g, 4.46 mmol) were added. The resultant solution was heated at reflux overnight under a nitrogen atmosphere. The following morning, the solvents were removed by rotary evaporation,  
20     and the residue was partitioned between  $\text{CH}_2\text{Cl}_2$  and saturated aqueous  $\text{NaHCO}_3$ . The organic phase was dried over  $\text{MgSO}_4$ , filtered, and concentrated to an orange oil. Purification via flash chromatography on silica gel (2% MeOH in  $\text{CH}_2\text{Cl}_2$  eluent) afforded 820 mg (46% yield) of 6-(2-ethoxyethyl)- $N^2,N^2$ -bis(4-methoxybenzyl)- $N^4$ -[4-(methylsulfonyl)butyl]-3-nitropyridine-2,4-diamine as a thick orange oil that was clean by  
25      $^1\text{H}$  NMR analysis.

Part K

To a solution of the material from Part J (820 mg, 1.37 mmol) in a 2:1  $\text{CH}_2\text{Cl}_2$  / MeOH mixture (50 mL) was added nickel(II) chloride hexahydrate (162 mg, 0.68 mmol) and  $\text{NaBH}_4$  (93 mg, 2.5 mmol). The solution instantly became black with some frothing.  
30     After 1 hour, the reaction solution was filtered through CELITE filter agent and the filter cake was washed with additional  $\text{CH}_2\text{Cl}_2$ . The filtrate was then washed with saturated aqueous  $\text{NaHCO}_3$  and brine, dried over  $\text{MgSO}_4$ , filtered, and concentrated to afford 760

mg (97% yield) of 6-(2-ethoxyethyl)-*N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-*N*<sup>4</sup>-[4-(methylsulfonyl)butyl]pyridine-2,3,4-triamine as a clear, colorless oil that <sup>1</sup>H NMR analysis showed to be of sufficient purity to carry forward without further purification.

#### Part L


5           To a solution of the material from Part K (760 mg, 1.33 mmol) in THF (50 mL) was added *N,N'*-carbonyl diimidazole (324 mg, 2.00 mmol). The resultant dark green solution was heated at reflux overnight under a nitrogen atmosphere. The following morning, solvents were removed by rotary evaporation, and the residue was purified via flash chromatography on silica gel (2% MeOH in CH<sub>2</sub>Cl<sub>2</sub> eluent) to provide 720 mg (91%  
10 yield) of 4-[bis(4-methoxybenzyl)amino]-6-(2-ethoxyethyl)-1-[4-(methylsulfonyl)butyl]-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a thick yellow oil that solidified upon standing. <sup>1</sup>H NMR analysis indicated very clean product.

#### Part M

          The material from Part L (720 mg, 1.21 mmol) was dissolved in TFA (15 mL), and  
15 the resultant deep violet solution was allowed to stir at room temperature overnight. The following morning, the TFA was removed via rotary evaporation, and the residue was diluted with de-ionized water. The pH was then adjusted to 8-9 by addition of Na<sub>2</sub>CO<sub>3</sub>, and the solution was extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 x 50 mL) and a 3:1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH mixture (60 mL). The combined organic layers were washed with brine, dried over MgSO<sub>4</sub>,  
20 filtered, and concentrated to a tan solid. Purification via flash column chromatography on silica gel (ramp eluent from 4-10% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) afforded 250 mg (58% yield) of 4-amino-6-(2-ethoxyethyl)-1-[4-(methylsulfonyl)butyl]-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a white solid, mp 203-206 °C. <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ 10.2 (s, 1H), 6.50 (s, 1H), 5.61 (s, 2H), 3.73 (m, 2H), 3.63 (t, *J* = 7.2 Hz, 2H), 3.42 (q, *J* = 7.0 Hz, 2H), 3.33 (s, 3H),  
25 3.16 (m, 2H), 2.93 (s, 3H), 2.74 (t, *J* = 7.2 Hz, 2H), 1.71 (m, 4H), 1.09 (t, *J* = 7.0 Hz, 3H). MS *m/z* 357 (*M* + *H*<sup>+</sup>); Anal. calcd for C<sub>15</sub>H<sub>24</sub>N<sub>4</sub>O<sub>4</sub>S: C, 50.54; H, 6.79; N, 15.72. Found: C, 50.47; H, 6.68; N, 15.57.

## Example 19

4-Amino-1-{{3-(4-fluorophenyl)isoxazol-5-yl}methyl}-6,7-dimethyl-  
1*H*-imidazo[4,5-*c*]pyridin-2-ol



## 5 Part A

To a solution of 2-chloro-5,6-dimethyl-3-nitro-*N*-prop-2-ynylpyridin-4-amine (10.0 g, 41.7 mmol), see International Publication No. WO2006/065280 (Moser *et al.*) Example 18, in toluene (200 mL) was added di-*para*-methoxy benzylamine (16.1 g, 62.6 mmol) and triethylamine (8.7 mL, 62.6 mmol). The resultant solution was heated to reflux for 58 hours. Upon cooling, solvents were removed by rotary evaporation, and the residue was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and saturated aqueous NaHCO<sub>3</sub>. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 100 mL), and the combined organic layers were washed with brine, dried over MgSO<sub>4</sub>, filtered, and concentrated to afford a waxy red solid. Purification via suction filter chromatography on silica gel (3/1 hexane / EtOAc eluent) afforded 18.5 g (96% yield) of *N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-5,6-dimethyl-3-nitro-*N*<sup>4</sup>-prop-2-ynylpyridine-2,4-diamine as a bright orange solid that was quite pure by <sup>1</sup>H NMR analysis.

## 15 Part B

*N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-5,6-dimethyl-3-nitro-*N*<sup>4</sup>-prop-2-ynylpyridine-2,4-diamine (11.5 g, 25.0 mmol) was dissolved in a 1:1 EtOH / CH<sub>3</sub>CN mixture (300 mL), and a solution of sodium dithionite (21.7 g, 125 mmol) in de-ionized water (100 mL) was added. A precipitate formed immediately, and the resultant mixture was allowed to stir at room temperature for 45 min. The precipitate was then removed by filtration through a pad of CELITE filter agent and the filter cake was washed with CH<sub>2</sub>Cl<sub>2</sub>. The volatile solvents were removed by rotary evaporation, and the remaining residue was partitioned between saturated aqueous NaHCO<sub>3</sub> and EtOAc. The aqueous phase was extracted with additional EtOAc (3 x 100 mL), and the combined organic layers were washed with brine,

dried over  $\text{MgSO}_4$ , filtered, and concentrated to afford 7.13 g (66% yield) of  $N^2, N^2$ -bis(4-methoxybenzyl)-5,6-dimethyl- $N^4$ -prop-2-ynylpyridine-2,3,4-triamine as a yellow solid. Analysis by  $^1\text{H}$  NMR revealed product of sufficient purity to carry forward without additional purification.

5 Step C

To a solution of the material from Part B (7.13 g, 16.6 mmol) in THF (100 mL) was added  $N, N'$ -carbonyl diimidazole (4.03 g, 24.8 mmol). The resultant solution was heated at reflux for 24 hours under a nitrogen atmosphere. Upon cooling, the solvents were removed by rotary evaporation, and the residue was partitioned between  $\text{CH}_2\text{Cl}_2$  and  
10 aqueous 1N HCl. The organic phase was subsequently washed with brine, dried over  $\text{MgSO}_4$ , filtered, and concentrated to an orange solid. Initial purification via suction filter chromatography on silica gel (1/1 EtOAc /  $\text{CH}_2\text{Cl}_2$  eluent) removed non-polar impurities; subsequent crystallization from EtOAc provided 1.5 g of 4-[bis(4-methoxybenzyl)amino]-6,7-dimethyl-1-prop-2-ynyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a white crystalline solid,  
15 along with an additional 5 g of material which required further purification.

Part D

A solution of 4-fluorobenzaldehyde oxime (0.97 g, 7.0 mmol), see International Publication No. WO2006/065280 (Moser *et al.*) Example 11, in DMF (14 mL) was chilled in an ice/water bath.  $N$ -Chlorosuccinimide (0.93 g, 7.0 mmol) was added in a single  
20 portion. The ice bath was removed; the solution was stirred at ambient temperature for 2 hours and then partitioned between ethyl acetate (50 mL) and water (50 mL). The layers were separated and the aqueous layer was extracted with ethyl acetate (2 x 50 mL). The combined organics were washed with brine (50 mL), dried over sodium sulfate, filtered, and then concentrated under reduced pressure to provide about 1.2 g of 4-fluoro- $N$ -  
25 hydroxybenzenecarboximidoyl chloride as a light yellow solid.

Part E

A suspension of the material from part D (about 7 mmol) in chloroform (23 mL) was cooled in an ice/water bath. Solid 4-[bis(4-methoxybenzyl)amino]-6,7-dimethyl-1-prop-2-ynyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol (1.3 g, 2.8 mmol) was added, followed by  
30 triethylamine (1.5 mL, 10.5 mmol). The ice bath was removed and the suspension was stirred at ambient temperature for 18 hours by which time a solution had been obtained. The reaction was quenched with saturated aqueous ammonium chloride (30 mL). The




organic layer was dried over sodium sulfate, filtered, and then concentrated under reduced pressure to provide an oil. This material was purified by automated flash chromatography (40+M cartridge, eluting with 20 to 60% ethyl acetate in hexanes) to provide 0.84 g of 4-[bis(4-methoxybenzyl)amino]-1-{[3-(4-fluorophenyl)isoxazol-5-yl]methyl}-6,7-dimethyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol.

## Part F

A solution of the material from Part E in trifluoroacetic acid (4 mL) was stirred at ambient temperature for 2 hours. The reaction was quenched with water (20 mL) and a white precipitate formed. The pH of the suspension was adjusted to about 13 with 50% aqueous sodium hydroxide. The mixture was stirred for 1 hour. The solid was isolated by filtration, washed with water, and then purified by automated flash chromatography (40+M cartridge, eluting with 0 to 15% methanol in dichloromethane) to provide a solid. The solid was combined with ethanol (60 mL) and heated to reflux. The mixture was allowed to cool. A solid was isolated by filtration, washed with ethanol, and then dried under vacuum at 65 °C to provide 0.3 g of 4-amino-1-{{[3-(4-fluorophenyl)]isoxazol-5-yl}methyl}-6,7-dimethyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a crystalline solid, mp 250.0 °C. Anal. Calcd for C<sub>18</sub>H<sub>16</sub>FN<sub>5</sub>O<sub>2</sub>: C, 61.18; H, 4.56; N, 19.82. Found: C, 60.99; H, 4.62; N, 19.55.

### Example 20

4-Amino-6,7-dimethyl-1-[(3-methylisoxazol-5-yl)methyl]-  
1*H*-imidazo[4,5-*c*]pyridin-2-ol



## Part A

25 A solution of acetaldoxime (0.43 g, 7.4 mmol) and *N*-chlorosuccinimide (0.98 g, 7.4 mmol) in DMF (22 mL) was heated at 50 °C for 2 hours. The reaction mixture was allowed to cool to ambient temperature; then it was quenched with water (150 mL) and extracted with ethyl acetate (2 x 50 mL). The combined organics were washed with brine

(50 mL), dried over sodium sulfate, filtered, and then concentrated under reduced pressure to provide 0.45 g of *N*-hydroxyethanimidoyl chloride as a clear oil.

#### Part B

5 A solution of the material from Part A (4.8 mmol) in dichloromethane (25 mL) was cooled in an ice/water bath. Solid 4-[bis(4-methoxybenzyl)amino]-6,7-dimethyl-1-prop-2-ynyl-1*H*-imidazo[4,5-*c*]pyridin-2-ol (1.8 g, 3.9 mmol) was added, followed by triethylamine (0.82 mL, 5.9 mmol). The ice bath was removed. The reaction solution was stirred at ambient temperature for 68 hours and then concentrated under reduced pressure to provide crude product as an oil. This material was purified by automated flash  
10 chromatography (40+M cartridge, eluting with 0% to 5% methanol in dichloromethane) to provide 0.58 g of 4-[bis(4-methoxybenzyl)amino]-6,7-dimethyl-1-[(3-methylisoxazol-5-yl)methyl]-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a white solid.


#### Part C

A solution of the material from Part B in trifluoroacetic acid (3 mL) was stirred at  
15 ambient temperature for 18 hours. The reaction was quenched with water (20 mL). The pH was adjusted to about 14 with 50% aqueous sodium hydroxide and then the mixture was neutralized (pH 7) with 1M aqueous hydrochloric acid. The resulting suspension was stirred for 1 hour. The solid was isolated by filtration, washed with water, and dried. This material was purified by automated flash chromatography (25+M cartridge, eluting with  
20 50% to 100% CMA in chloroform) followed by recrystallization from methanol to provide 0.13 g of 4-amino-6,7-dimethyl-1-[(3-methylisoxazol-5-yl)methyl]-1*H*-imidazo[4,5-*c*]pyridin-2-ol as a crystalline solid, mp >300 °C. Anal. Calcd for C<sub>13</sub>H<sub>15</sub>N<sub>5</sub>O<sub>2</sub>: C, 57.13; H, 5.53; N, 25.63. Found: C, 57.15; H, 5.75; N, 25.81.

25

#### Example 21

4-Amino-6-pentyl-1-[(1*S*)-1-phenylethyl]-1*H*-imidazo[4,5-*c*]pyridin-2-ol



## Part A

A solution of 2,4-dichloro-3-nitro-6-pentylpyridine (1.11 g, 4.22 mmol) dissolved in 10 mL of *N,N*-dimethylformamide was treated with triethylamine (1.17 mL, 8.44 mmol) and (D)-(+)-phenethylamine (536 mL, 4.22 mmol). After stirring for 6 hours at ambient temperature, the reaction mixture was heated to 40 °C for 1 hour and then concentrated under reduced pressure. The residue was dissolved in 50 mL of ethyl acetate and washed successively with H<sub>2</sub>O (3x) and brine. The organic portion was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to give yellow oil. Column chromatography (SiO<sub>2</sub>, 3-15% methyl *tert*-butylether (MTBE)/hexanes) gave the 2-chloro-3-nitro-6-pentyl-*N*-[(1*S*)-1-phenylethyl]pyridin-4-amine (715 mg) as a yellow syrup.

## Part B

A solution of 2-chloro-3-nitro-6-pentyl-*N*-[(1*S*)-1-phenylethyl]pyridin-4-amine (715 mg, 2.05 mmol) dissolved in 20 mL of toluene was treated with triethylamine (0.57 mL, 4.1 mmol) and di-*p*-methoxybenzylamine (581 mg, 2.26 mmol) and the mixture was heated to reflux overnight. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in 50 mL of ethyl acetate and washed successively with H<sub>2</sub>O and brine. The organic portion was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to give brown oil. Column chromatography (SiO<sub>2</sub>, 3-15% MTBE/hexanes) gave the *N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-3-nitro-6-pentyl-*N*<sup>4</sup>-[(1*S*)-1-phenylethyl]pyridine-2,4-diamine (1.10 g) as a dark-yellow syrup.

## Part C

A stirred solution of Ni(II)chloride hexahydrate (230 mg) dissolved in 10 mL of methanol was treated with NaBH<sub>4</sub> (74 mg). A solution of *N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-3-nitro-6-pentyl-*N*<sup>4</sup>-[(1*S*)-1-phenylethyl]pyridine-2,4-diamine (1.10 g, 1.94 mmol), dissolved in 10 mL of a 1:1 mixture of methanol/CH<sub>2</sub>Cl<sub>2</sub>, was then added to the stirred solution. Additional 20 mg-portions of NaBH<sub>4</sub> (about 8) were then added over 30 minutes until the reaction mixture turned from yellow-brown to clear. Thin-layer chromatography showed the complete consumption of starting material. The reaction mixture was then filtered through a layer of CELITE filter agent. The filter cake was rinsed with additional CH<sub>2</sub>Cl<sub>2</sub> and the combined filtrates were concentrated under reduced pressure. The resulting material was then filtered through a short column of SiO<sub>2</sub>, eluting with 5-10%

methanol/CHCl<sub>3</sub> to give *N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-6-pentyl-*N*<sup>4</sup>-[(1*S*)-1-phenylethyl]pyridine-2,3,4-triamine (979 mg) as a light brown solid.

#### Part D

A solution of *N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-6-pentyl-*N*<sup>4</sup>-[(1*S*)-1-phenylethyl]pyridine-2,3,4-triamine (979 mg, 1.82 mmol) dissolved in 10 mL of THF was treated with carbonyl diimidazole (590 mg, 3.64 mmol) and the mixture was heated to reflux. After 90 minutes, the reaction mixture was treated with an additional portion (200 mg) of carbonyl diimidazole and heating was continued for 1 hour. The reaction mixture was cooled and treated with 10 mL of H<sub>2</sub>O. After stirring for 10 minutes, the reaction mixture was diluted with 30 mL of ethyl acetate. The layers were separated and the organic portion was washed successively with H<sub>2</sub>O and brine. The organic portion was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to give a purple oil. Column chromatography (SiO<sub>2</sub>, 0-5% methanol/CH<sub>2</sub>Cl<sub>2</sub>) gave the 4-[bis(4-methoxybenzyl)amino]-6-pentyl-1-[(1*S*)-1-phenylethyl]-1*H*-imidazo[4,5-*c*]pyridin-2-ol (977 mg) as a purple syrup.

#### Part E


A solution of 4-[bis(4-methoxybenzyl)amino]-6-pentyl-1-[(1*S*)-1-phenylethyl]-1*H*-imidazo[4,5-*c*]pyridin-2-ol (977 mg, 1.73 mmol) dissolved in 10 mL of trifluoroacetic acid (TFA) was stirred overnight. The reaction mixture was then concentrated under reduced pressure and the resulting residue was partitioned between dilute NH<sub>4</sub>OH and CH<sub>2</sub>Cl<sub>2</sub>. The layers were separated and the aqueous portion was extracted with 2 additional portions of CH<sub>2</sub>Cl<sub>2</sub>. The combined organic portions were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated under reduced pressure. Column chromatography (SiO<sub>2</sub>, 10-35% CMA/CHCl<sub>3</sub>) gave the title compound, which was further purified by crystallization from 10 mL of hot acetonitrile. The crystals were isolated by filtration and dried under high vacuum to give 4-amino-6-pentyl-1-[(1*S*)-1-phenylethyl]-1*H*-imidazo[4,5-*c*]pyridin-2-ol (340 mg) as pinkish crystals: mp 146.7-147.7 °C; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 10.90 (br s, 1H), 7.22-7.39 (m, 5H), 6.02 (s, 1H), 5.74 (q, *J* = 7.2 Hz, 1H), 4.92 (s, 2H), 2.48 (t, *J* = 7.7 Hz, 2H), 1.89 (d, *J* = 7.2 Hz, 1H), 1.53 (m, 2H), 1.17-1.29 (m, 4H), 0.84 (t, *J* = 7.1 Hz, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 154.9, 154.0, 142.7, 139.4, 136.0, 128.7, 127.7, 126.4, 108.4, 96.5, 51.0, 38.4, 31.5, 29.8, 22.5, 17.8, 14.0; MS *m/z* 325 (M + H)<sup>+</sup>. Anal.

calcd for C<sub>19</sub>H<sub>24</sub>N<sub>4</sub>O: C, 70.34; H, 7.46; N, 17.27. Found: C, 70.30; H, 7.51; N, 17.22.

Optical rotation [ $\alpha$ ]=-58.6 (c=1.33 mg/mL, CH<sub>2</sub>Cl<sub>2</sub>).

#### Example 22

5            4-Amino-6-pentyl-1-[(1*R*)-1-phenylethyl]-1*H*-imidazo[4,5-*c*]pyridin-2-ol




4-Amino-6-pentyl-1-[(1*R*)-1-phenylethyl]-1*H*-imidazo[4,5-*c*]pyridin-2-ol was prepared according to the methods described in Parts A through E of Example 21 using (L)-(-)-phenethylamine in lieu of (D)-(+)-phenethylamine in Part A. mp 147.1-148.0 °C;

10        Anal. calcd for C<sub>19</sub>H<sub>24</sub>N<sub>4</sub>O: C, 70.34; H, 7.46; N, 17.27. Found: C, 70.41; H, 7.51; N, 17.40. Optical rotation [ $\alpha$ ]=+61.5 (c=1.82 mg/mL, CH<sub>2</sub>Cl<sub>2</sub>).

#### Example 23

4-amino-1-benzyl-6-butoxy-1*H*-imidazo[4,5-*c*]pyridin-2-ol



15

#### Part A

4-Amino-2,6-dichloropyridine (1.30 g, 7.98 mmol) was carefully added to 6.5 mL of concentrated sulfuric acid. The mixture was cooled in an ice bath, and 2.6 mL of fuming nitric acid was added dropwise via pipette. The solution was warmed to room temperature, stirred for one hour, and then poured onto 26 g of crushed ice, resulting in the formation of a white precipitate. The mixture was stored at -10 °C overnight. The white precipitate was collected by filtration using a Buchner funnel, washed with ice cold water, and dried under vacuum to provide 1.66 g of 2,6-dichloro-4-nitraminopyridine, which was carried forward without additional purification.

20

## Part B

2,6-Dichloro-4-nitraminopyridine (1.66 g, 7.98 mmol) was added to 11 mL of concentrated sulfuric acid, and the resultant solution was heated on a steam bath for 30 minutes. After cooling to room temperature, the solution was poured onto 28 g of crushed ice, resulting in the formation of a tan precipitate. The mixture was cooled in an ice bath, and concentrated ammonium hydroxide was added until pH 7 was reached. The resultant slurry was stored at -10 °C overnight. The precipitate was collected by filtration using a Buchner funnel, washed with ice cold water, and dried under vacuum to provide 4-amino-2,6-dichloro-3-nitropyridine (1.30 g, 78% yield) as a light tan solid.

## Part C

A solution of 4-amino-2,6-dichloro-3-nitropyridine (1.08 g, 5.19 mmol) from Part B and triethylamine (1.09 mL, 7.79 mmol) in dichloromethane (20 mL) was cooled in an ice bath, and bis(4-methoxybenzyl)amine (1.34 g, 5.19 mmol) was added in one portion. The resultant solution was allowed to warm to room temperature and stirred overnight under a nitrogen atmosphere. The solvents were removed by rotary evaporation, and the residue was purified via flash chromatography (silica gel, dichloromethane eluent) to provide *N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-6-chloro-3-nitropyridine-2,4-diamine (1.82 g, 82% yield) as a viscous yellow oil that foamed under vacuum.

## Part D

Sodium butoxide was prepared by adding sodium metal (207 mg, 9.00 mmol) to 1-butanol (7 mL). After the sodium metal had been completely consumed, the resulting solution was cooled in an ice bath, and a solution of *N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-6-chloro-3-nitropyridine-2,4-diamine (1.29 g, 3.00 mmol) from Part C in THF (10 mL) was added dropwise via an addition funnel. The resultant solution was heated in an 85 °C oil bath for five hours. Upon cooling to room temperature, the reaction mixture was quenched by the addition of dilute aqueous HCl and extracted with dichloromethane (3 x 50 mL). The combined organic layers were washed with brine, dried over MgSO<sub>4</sub>, filtered, and concentrated to a red oil. Purification via flash chromatography (silica gel, dichloromethane eluent) provided *N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-6-butoxy-3-nitropyridine-2,4-diamine (1.14 g, 81% yield) as a viscous orange oil that foamed under vacuum.

## Part E

*N*<sup>2</sup>,*N*<sup>2</sup>-Bis(4-methoxybenzyl)-6-butoxy-3-nitropyridine-2,4-diamine (1.10 g, 2.36 mmol) from Part D was dissolved in 40 mL of a 1:1 ethanol/acetonitrile mixture, and a solution of sodium hydrosulfite, Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub>, (2.05 g, 11.8 mmol) in H<sub>2</sub>O (10 mL) was added via pipette, resulting in the formation of a white precipitate. The mixture was stirred at room temperature for two hours, during which time the orange-yellow color faded away. The mixture was then filtered through a pad of CELITE filter agent, the filter cake was washed with dichloromethane, and the filtrate was concentrated under reduced pressure. The residue was diluted with ethyl acetate (150 mL), washed with saturated aqueous sodium bicarbonate (1 x 50 mL) and brine (1 x 25 mL), dried over magnesium sulfate, filtered, and concentrated to provide *N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-6-butoxy-pyridine-2,3,4-diamine as a yellow oil. This material was carried forward without additional purification.

## Part F

To a solution of *N*<sup>2</sup>,*N*<sup>2</sup>-bis(4-methoxybenzyl)-6-butoxy-pyridine-2,3,4-diamine (1.03 g, 2.35 mmol) from Part E in THF (25 mL) was added 1,1'-carbonyldiimidazole (496 mg, 3.06 mmol). The resultant solution was heated to reflux overnight under a nitrogen atmosphere. The solvents were removed by rotary evaporation, and the residue was purified via flash chromatography (silica gel, ramp eluent from 2/1 to 1/2 hexane/ethyl acetate) to provide 6-butoxy-4-[di(4-methoxybenzyl)amino]-1,3-dihydroimidazo[4,5-*c*]pyridin-2-one (630 mg, 58% yield over two steps) as a light pink oil that foamed under vacuum.

## Part G

The 6-butoxy-4-[di(4-methoxybenzyl)amino]-1,3-dihydroimidazo[4,5-*c*]pyridin-2-one (630 mg, 1.36 mmol) from Part F was dissolved in DMF (8 mL), and solid potassium carbonate (225 mg, 1.63 mmol) was added. A solution of benzyl bromide (257 mg, 1.50 mmol) in DMF (2 mL) was then added via pipette, and the resultant solution was heated at 80 °C in an oil bath overnight under a nitrogen atmosphere. The reaction mixture was diluted with ethyl acetate (150 mL), washed with water (4 x 25 mL) and brine (4 x 25 mL), dried over magnesium sulfate, filtered, and concentrated to give an oil. Purification by flash chromatography (silica gel, ramp eluent from 2/1 to 1/2 hexane/ethyl acetate)


provided 1-benzyl-6-butoxy-4-[di(4-methoxybenzyl)amino]-1*H*-imidazo[4,5-*c*]pyridin-2-ol (80 mg, 11% yield) as a tan oil.

#### Part H

To the 1-benzyl-6-butoxy-4-[di(4-methoxybenzyl)amino]-1*H*-imidazo[4,5-*c*]pyridin-2-ol (80 mg, 0.15 mmol) from Part G was added TFA (5 mL), forming a deep violet solution, which was allowed to stir at room temperature overnight. The TFA was removed via rotary evaporation, and the residue was diluted with water (10 mL). Solid sodium carbonate was added to adjust the pH to about 8-9. The aqueous layer was extracted with dichloromethane (3 x 25 mL). The combined organic layers were dried over magnesium sulfate, filtered, and concentrated to give a tan solid. The tan solid was purified by flash chromatography (silica gel, 6% methanol in dichloromethane eluent) to provide 4-amino-1-benzyl-6-butoxy-1*H*-imidazo[4,5-*c*]pyridin-2-ol (30 mg, 66% yield) as a light tan solid, mp 205-208 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 9.41 (s, 1H), 7.47-7.24 (m, 5H), 5.99 (s, 1H), 5.22 (s, 2H), 4.08 (t, *J* = 7.1 Hz, 2H), 3.98 (s, 2H), 1.70 (m, 2H), 1.44 (m, 2H), 0.94 (t, *J* = 7.1 Hz, 3H). MS *m/z* 313 (*M* + H<sup>+</sup>).

#### Exemplary Compounds

Certain exemplary compounds, including some of those described above in the Examples, have the following Formula (Ia) and an R<sub>1</sub> substituent shown in the following table, wherein each line of the table is matched with the Formula (Ia) to represent a specific embodiment of the invention.



Ia

R <sub>1</sub>
pyridin-3-ylmethyl
4-fluorobenzyl
isoxazol-5-ylmethyl
isoxazol-3-ylmethyl



[5-(4-fluorophenyl)isoxazol-3- yl]methyl
[3-(4-fluorophenyl)isoxazol-5- yl]methyl

Compounds of the invention have been found to modulate cytokine biosynthesis by inducing the production of interferon  $\alpha$ , or interferon  $\alpha$  and tumor necrosis factor  $\alpha$  in human cells when tested using one of the methods described below.

#### CYTOKINE INDUCTION IN HUMAN CELLS

An in vitro human blood cell system is used to assess cytokine induction. Activity is based on the measurement of interferon ( $\alpha$ ) and tumor necrosis factor ( $\alpha$ ) (IFN- $\alpha$  and TNF- $\alpha$ , respectively) secreted into culture media as described by Testerman et al. in "Cytokine Induction by the Immunomodulators Imiquimod and S-27609", *Journal of Leukocyte Biology*, 58, 365-372 (September, 1995).

##### Blood Cell Preparation for Culture

Whole blood from healthy human donors is collected by venipuncture into vacutainer tubes or syringes containing EDTA. Peripheral blood mononuclear cells (PBMC) are separated from whole blood by density gradient centrifugation using HISTOPAQUE-1077 (Sigma, St. Louis, MO) or Ficoll-Paque Plus (Amersham Biosciences Piscataway, NJ). Blood is diluted 1:1 with Dulbecco's Phosphate Buffered Saline (DPBS) or Hank's Balanced Salts Solution (HBSS). Alternately, whole blood is placed in Accuspin (Sigma) or LeucoSep (Greiner Bio-One, Inc., Longwood, FL) centrifuge frit tubes containing density gradient medium. The PBMC layer is collected and washed twice with DPBS or HBSS and re-suspended at  $4 \times 10^6$  cells/mL in RPMI complete. The PBMC suspension is added to 96 well flat bottom sterile tissue culture plates containing an equal volume of RPMI complete media containing test compound.

### Compound Preparation

The compounds are solubilized in dimethyl sulfoxide (DMSO). The DMSO concentration should not exceed a final concentration of 1% for addition to the culture wells. The compounds are generally tested at concentrations ranging from 30-0.014  $\mu\text{M}$ . Controls include cell samples with media only, cell samples with DMSO only (no compound), and cell samples with reference compound.

### Incubation

The solution of test compound is added at 60  $\mu\text{M}$  to the first well containing RPMI complete and serial 3 fold dilutions are made in the wells. The PBMC suspension is then added to the wells in an equal volume, bringing the test compound concentrations to the desired range (usually 30-0.014  $\mu\text{M}$ ). The final concentration of PBMC suspension is  $2 \times 10^6$  cells/mL. The plates are covered with sterile plastic lids, mixed gently and then incubated for 18 to 24 hours at 37°C in a 5% carbon dioxide atmosphere.

### Separation

Following incubation the plates are centrifuged for 10 minutes at 1000 rpm (approximately 200 x g) at 4°C. The cell-free culture supernatant is removed and transferred to sterile polypropylene tubes. Samples are maintained at -30 to -70°C until analysis. The samples are analyzed for IFN- $\alpha$  by ELISA and for TNF- $\alpha$  by IGEN/BioVeris Assay.

### Interferon ( $\alpha$ ) and Tumor Necrosis Factor ( $\alpha$ ) Analysis

IFN- $\alpha$  concentration is determined with a human multi-subtype colorimetric sandwich ELISA (Catalog Number 41105) from PBL Biomedical Laboratories, Piscataway, NJ. Results are expressed in pg/mL.

The TNF- $\alpha$  concentration is determined by ORIGEN M-Series Immunoassay and read on an IGEN M-8 analyzer from BioVeris Corporation, formerly known as IGEN International, Gaithersburg, MD. The immunoassay uses a human TNF- $\alpha$  capture and detection antibody pair (Catalog Numbers AHC3419 and AHC3712) from Biosource International, Camarillo, CA. Results are expressed in pg/mL.

### Assay Data and Analysis

In total, the data output of the assay consists of concentration values of TNF- $\alpha$  and IFN- $\alpha$  (y-axis) as a function of compound concentration (x-axis).

Analysis of the data has two steps. First, the greater of the mean DMSO (DMSO control wells) or the experimental background (usually 20 pg/mL for IFN- $\alpha$  and 40 pg/mL for TNF- $\alpha$ ) is subtracted from each reading. If any negative values result from background subtraction, the reading is reported as " \* ", and is noted as not reliably detectable. In subsequent calculations and statistics, " \* ", is treated as a zero. Second, all background subtracted values are multiplied by a single adjustment ratio to decrease experiment to experiment variability. The adjustment ratio is the area of the reference compound in the new experiment divided by the expected area of the reference compound based on the past 61 experiments (unadjusted readings). This results in the scaling of the reading (y-axis) for the new data without changing the shape of the dose-response curve. The reference compound used is 2-[4-amino-2-ethoxymethyl-6,7,8,9-tetrahydro- $\alpha$ , $\alpha$ -dimethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl]ethanol hydrate (U.S. Patent No. 5,352,784; Example 91) and the expected area is the sum of the median dose values from the past 61 experiments.

The minimum effective concentration is calculated based on the background-subtracted, reference-adjusted results for a given experiment and compound. The minimum effective concentration ( $\mu$ molar) is the lowest of the tested compound concentrations that induces a response over a fixed cytokine concentration for the tested cytokine (usually 20 pg/mL for IFN- $\alpha$  and 40 pg/mL for TNF- $\alpha$ ). The maximal response is the maximal amount of cytokine (pg/ml) produced in the dose-response.

### CYTOKINE INDUCTION IN HUMAN CELLS (High Throughput Screen)

The CYTOKINE INDUCTION IN HUMAN CELLS test method described above was modified as follows for high throughput screening.

#### Blood Cell Preparation for Culture

Whole blood from healthy human donors is collected by venipuncture into vacutainer tubes or syringes containing EDTA. Peripheral blood mononuclear cells

(PBMC) are separated from whole blood by density gradient centrifugation using HISTOPAQUE-1077 (Sigma, St. Louis, MO) or Ficoll-Paque Plus (Amersham Biosciences Piscataway, NJ). Whole blood is placed in Accuspin (Sigma) or LeucoSep (Greiner Bio-One, Inc., Longwood, FL) centrifuge frit tubes containing density gradient medium. The PBMC layer is collected and washed twice with DPBS or HBSS and re-suspended at  $4 \times 10^6$  cells/mL in RPMI complete (2-fold the final cell density). The PBMC suspension is added to 96-well flat bottom sterile tissue culture plates.

#### Compound Preparation

The compounds are solubilized in dimethyl sulfoxide (DMSO). The compounds are generally tested at concentrations ranging from 30 - 0.014  $\mu$ M. Controls include cell samples with media only, cell samples with DMSO only (no compound), and cell samples with a reference compound 2-[4-amino-2-ethoxymethyl-6,7,8,9-tetrahydro- $\alpha,\alpha$ -dimethyl-1H-imidazo[4,5-c]quinolin-1-yl]ethanol hydrate (U.S. Patent No. 5,352,784; Example 91) on each plate. The solution of test compound is added at 7.5 mM to the first well of a dosing plate and serial 3 fold dilutions are made for the 7 subsequent concentrations in DMSO. RPMI Complete media is then added to the test compound dilutions in order to reach a final compound concentration of 2-fold higher (60 - 0.028  $\mu$ M) than the final tested concentration range.

#### Incubation

Compound solution is then added to the wells containing the PBMC suspension bringing the test compound concentrations to the desired range (usually 30 - 0.014  $\mu$ M) and the DMSO concentration to 0.4 %. The final concentration of PBMC suspension is  $2 \times 10^6$  cells/mL. The plates are covered with sterile plastic lids, mixed gently and then incubated for 18 to 24 hours at 37°C in a 5% carbon dioxide atmosphere.

#### Separation

Following incubation the plates are centrifuged for 10 minutes at 1000 rpm (approximately 200 g) at 4°C. 4-plex Human Panel MSD MULTI-SPOT 96-well plates are pre-coated with the appropriate capture antibodies by MesoScale Discovery, Inc. (MSD, Gaithersburg, MD). The cell-free culture supernatants are removed and transferred

to the MSD plates. Fresh samples are typically tested, although they may be maintained at -30 to -70°C until analysis.

#### Interferon- $\alpha$ and Tumor Necrosis Factor- $\alpha$ Analysis

5 MSD MULTI-SPOT plates contain within each well capture antibodies for human TNF- $\alpha$  and human IFN- $\alpha$  that have been pre-coated on specific spots. Each well contains four spots: one human TNF- $\alpha$  capture antibody (MSD) spot, one human IFN- $\alpha$  capture antibody (PBL Biomedical Laboratories, Piscataway, NJ) spot, and two inactive bovine serum albumin spots. The human TNF- $\alpha$  capture and detection antibody pair is from  
10 MesoScale Discovery. The human IFN- $\alpha$  multi-subtype antibody (PBL Biomedical Laboratories) captures all IFN- $\alpha$  subtypes except IFN- $\alpha$  F (IFNA21). Standards consist of recombinant human TNF- $\alpha$  (R&D Systems, Minneapolis, MN) and IFN- $\alpha$  (PBL Biomedical Laboratories). Samples and separate standards are added at the time of analysis to each MSD plate. Two human IFN- $\alpha$  detection antibodies (Cat. Nos. 21112 &  
15 21100, PBL) are used in a two to one ratio (weight:weight) to each other to determine the IFN- $\alpha$  concentrations. The cytokine-specific detection antibodies are labeled with the SULFO-TAG reagent (MSD). After adding the SULFO-TAG labeled detection antibodies to the wells, each well's electrochemoluminescent levels are read using MSD's SECTOR HTS READER. Results are expressed in pg/mL upon calculation with known cytokine  
20 standards.

#### Assay Data and Analysis

In total, the data output of the assay consists of concentration values of TNF- $\alpha$  or IFN- $\alpha$  (y-axis) as a function of compound concentration (x-axis).

25 A plate-wise scaling is performed within a given experiment aimed at reducing plate-to-plate variability associated within the same experiment. First, the greater of the median DMSO (DMSO control wells) or the experimental background (usually 20 pg/mL for IFN- $\alpha$  and 40 pg/mL for TNF- $\alpha$ ) is subtracted from each reading. Negative values that may result from background subtraction are set to zero. Each plate within a given  
30 experiment has a reference compound that serves as a control. This control is used to calculate a median expected area under the curve across all plates in the assay. A plate-wise scaling factor is calculated for each plate as a ratio of the area of the reference

compound on the particular plate to the median expected area for the entire experiment. The data from each plate are then multiplied by the plate-wise scaling factor for all plates. Only data from plates bearing a scaling factor of between 0.5 and 2.0 (for both cytokines IFN- $\alpha$ , TNF- $\alpha$ ) are reported. Data from plates with scaling factors outside the above mentioned interval are retested until they bear scaling factors inside the above mentioned interval. The above method produces a scaling of the y-values without altering the shape of the curve. The reference compound used is 2-[4-amino-2-ethoxymethyl-6,7,8,9-tetrahydro- $\alpha,\alpha$ -dimethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl]ethanol hydrate (U.S. Patent No. 5,352,784; Example 91). The median expected area is the median area across all plates that are part of a given experiment.

A second scaling may also be performed to reduce inter-experiment variability (across multiple experiments). All background-subtracted values are multiplied by a single adjustment ratio to decrease experiment-to-experiment variability. The adjustment ratio is the area of the reference compound in the new experiment divided by the expected area of the reference compound based on an average of previous experiments (unadjusted readings). This results in the scaling of the reading (y-axis) for the new data without changing the shape of the dose-response curve. The reference compound used is 2-[4-amino-2-ethoxymethyl-6,7,8,9-tetrahydro- $\alpha,\alpha$ -dimethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl]ethanol hydrate (U.S. Patent No. 5,352,784; Example 91) and the expected area is the sum of the median dose values from an average of previous experiments.


The minimum effective concentration is calculated based on the background-subtracted, reference-adjusted results for a given experiment and compound. The minimum effective concentration ( $\mu$ molar) is the lowest of the tested compound concentrations that induces a response over a fixed cytokine concentration for the tested cytokine (usually 20 pg/mL for IFN- $\alpha$  and 40 pg/mL for TNF- $\alpha$ ). The maximal response is the maximal amount of cytokine (pg/ml) produced in the dose-response.

The complete disclosures of the patents, patent documents, and publications cited herein are incorporated by reference in their entirety as if each were individually incorporated. Various modifications and alterations to this invention will become apparent to those skilled in the art without departing from the scope and spirit of this invention. It should be understood that this invention is not intended to be unduly limited

by the illustrative embodiments and examples set forth herein and that such examples and embodiments are presented by way of example only with the scope of the invention intended to be limited only by the claims set forth herein as follows.

## WHAT IS CLAIMED IS:

1. A compound of the Formula I:



I

wherein:

$R_A$  and  $R_B$  are each independently selected from the group consisting of:

hydrogen,

halogen,

alkenyl,

amino,

$-R_{11}$ ,

$-O-R_{11}$ ,

$-S-R_{11}$ , and

$-N(R_{9a})(R_{11})$ ;

$R_{11}$  is selected from the group consisting of alkyl, alkoxyalkylenyl,

hydroxyalkylenyl, aryl, arylalkylenyl, heteroaryl, heteroarylalkylenyl, heterocyclyl, and

heterocyclylalkylenyl, each of which is unsubstituted or substituted by one or more

substituents independently selected from the group consisting of alkyl; alkoxy; hydroxy;

hydroxyalkyl; aryl; aryloxy; arylalkyleneoxy; heteroaryl; heteroaryloxy;

heteroarylalkyleneoxy; halogen; haloalkyl; haloalkoxy; mercapto; nitro; cyano;

heterocyclyl; amino; alkylamino; dialkylamino; and, in the case of alkyl, heterocyclyl, and

heterocyclylalkylenyl, oxo;

$R_{9a}$  is selected from the group consisting of hydrogen and  $C_{1-4}$  alkyl;

$R_1$  is selected from the group consisting of:

$-R_4$ ,

$-X-R_4$ ,

$-X-Y-R_4$ ,

$-X-Y-X-Y-R_4$ ,



-X-R<sub>5</sub>,  
-N(R<sub>1</sub>')-Q-R<sub>4</sub>,  
-N(R<sub>1</sub>')-X<sub>1</sub>-Y<sub>1</sub>-R<sub>4</sub>, and  
-N(R<sub>1</sub>')-X<sub>1</sub>-R<sub>5a</sub>;


5 X is selected from the group consisting of alkylene, alkenylene, alkynylene, arylene, heteroarylene, and heterocyclylene wherein the alkylene, alkenylene, and alkynylene groups can be optionally interrupted or terminated by arylene, heteroarylene or heterocyclylene and optionally interrupted by one or more -O- groups;



X<sub>1</sub> is C<sub>2-20</sub> alkylene;

10 Y is selected from the group consisting of:


-O-,  
-S(O)<sub>0-2</sub>-,  
-S(O)<sub>2</sub>-N(R<sub>8</sub>)-,  
-C(R<sub>6</sub>)-,  
-C(R<sub>6</sub>)-O-,  
-O-C(R<sub>6</sub>)-,  
-O-C(O)-O-,  
-N(R<sub>8</sub>)-Q-,  
-C(R<sub>6</sub>)-N(R<sub>8</sub>)-,  
-O-C(R<sub>6</sub>)-N(R<sub>8</sub>)-,  
-C(R<sub>6</sub>)-N(OR<sub>9</sub>)-,  
-O-N(R<sub>8</sub>)-Q-,  
-O-N=C(R<sub>4</sub>)-,  
-C(=N-O-R<sub>8</sub>)-,  
-CH(-N(-O-R<sub>8</sub>)-Q-R<sub>4</sub>)-,

25





Y<sub>1</sub> is selected from the group consisting of -O-, -S(O)<sub>0-2</sub>-, -S(O)<sub>2</sub>-N(R<sub>8</sub>)-,




5 R<sub>1</sub>' is selected from the group consisting of hydrogen, C<sub>1-20</sub> alkyl, hydroxy-C<sub>2-20</sub> alkylenyl, and alkoxy-C<sub>2-20</sub> alkylenyl;

R<sub>4</sub> is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl, heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl wherein the alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl, heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl groups can be unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl; alkoxy; hydroxyalkyl; haloalkyl; haloalkoxy; halogen; nitro; hydroxy; mercapto; cyano; aryl; aryloxy; arylalkyleneoxy; heteroaryl; heteroaryloxy; heteroarylalkyleneoxy; heterocyclyl; amino; alkylamino; dialkylamino; (dialkylamino)alkyleneoxy; and, in the case of alkyl, alkenyl, alkynyl, and heterocyclyl, oxo;


10

15

R<sub>5</sub> is selected from the group consisting of:



R<sub>5a</sub> is selected from the group consisting of:



R<sub>6</sub> is selected from the group consisting of =O and =S;

R<sub>7</sub> is C<sub>2-7</sub> alkylene;

5 R<sub>8</sub> is selected from the group consisting of hydrogen, C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, hydroxy-C<sub>1-10</sub> alkylenyl, C<sub>1-10</sub> alkoxy-C<sub>1-10</sub> alkylenyl, aryl-C<sub>1-10</sub> alkylenyl, and heteroaryl-C<sub>1-10</sub> alkylenyl;

R<sub>9</sub> is selected from the group consisting of hydrogen and alkyl;

R<sub>10</sub> is C<sub>3-8</sub> alkylene;

A is selected from the group consisting of -CH<sub>2</sub>-, -O-, -C(O)-, -S(O)<sub>0-2</sub>-, and  
10 -N(-Q-R<sub>4</sub>)-;

A' is selected from the group consisting of -O-, -S(O)<sub>0-2</sub>-, -N(-Q-R<sub>4</sub>)-, and -CH<sub>2</sub>-;

Q is selected from the group consisting of a bond, -C(R<sub>6</sub>)-, -C(R<sub>6</sub>)-C(R<sub>6</sub>)-, -S(O)<sub>2</sub>-, -C(R<sub>6</sub>)-N(R<sub>8</sub>)-W-, -S(O)<sub>2</sub>-N(R<sub>8</sub>)-, -C(R<sub>6</sub>)-O-, -C(R<sub>6</sub>)-S-, and -C(R<sub>6</sub>)-N(OR<sub>9</sub>)-;


V is selected from the group consisting of -C(R<sub>6</sub>)-, -O-C(R<sub>6</sub>)-, -N(R<sub>8</sub>)-C(R<sub>6</sub>)-, and  
15 -S(O)<sub>2</sub>-;

W is selected from the group consisting of a bond, -C(O)-, and -S(O)<sub>2</sub>-; and

a and b are independently integers from 1 to 6 with the proviso that a + b is ≤ 7;

or a pharmaceutically acceptable salt thereof.

20 2. A compound of the Formula II:



II

wherein:

G<sub>1</sub> is selected from the group consisting of:

25 -C(O)-R',  
α-aminoacyl,  
α-aminoacyl-α-aminoacyl,

- 5  
 -C(O)-O-R',  
 -C(O)-N(R'')R',  
 -C(=NY')-R',  
 -CH(OH)-C(O)-OY',  
 -CH(OC<sub>1-4</sub> alkyl)Y<sub>0</sub>,  
 -CH<sub>2</sub>Y<sub>2</sub>, and  
 -CH(CH<sub>3</sub>)Y<sub>2</sub>;

10 R' and R'' are independently selected from the group consisting of C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, phenyl, and benzyl, each of which may be unsubstituted or substituted by one or more substituents independently selected from the group consisting of halogen, hydroxy, nitro, cyano, carboxy, C<sub>1-6</sub> alkyl, C<sub>1-4</sub> alkoxy, aryl, heteroaryl, aryl-C<sub>1-4</sub> alkylenyl, heteroaryl-C<sub>1-4</sub> alkylenyl, halo-C<sub>1-4</sub> alkylenyl, halo-C<sub>1-4</sub> alkoxy, -O-C(O)-CH<sub>3</sub>, -C(O)-O-CH<sub>3</sub>, -C(O)-NH<sub>2</sub>, -O-CH<sub>2</sub>-C(O)-NH<sub>2</sub>, -NH<sub>2</sub>, and -S(O)<sub>2</sub>-NH<sub>2</sub>, with the proviso that R'' can also be hydrogen;

15  $\alpha$ -aminoacyl is an  $\alpha$ -aminoacyl group derived from an amino acid selected from the group consisting of racemic, D-, and L-amino acids;

Y' is selected from the group consisting of hydrogen, C<sub>1-6</sub> alkyl, and benzyl;

20 Y<sub>0</sub> is selected from the group consisting of C<sub>1-6</sub> alkyl, carboxy-C<sub>1-6</sub> alkylenyl, amino-C<sub>1-4</sub> alkylenyl, mono-N-C<sub>1-6</sub> alkylamino-C<sub>1-4</sub> alkylenyl, and di-N,N-C<sub>1-6</sub> alkylamino-C<sub>1-4</sub> alkylenyl;

Y<sub>2</sub> is selected from the group consisting of mono-N-C<sub>1-6</sub> alkylamino, di-N,N-C<sub>1-6</sub> alkylamino, morpholin-4-yl, piperidin-1-yl, pyrrolidin-1-yl, and 4-C<sub>1-4</sub> alkylpiperazin-1-yl;

25 R<sub>A</sub> and R<sub>B</sub> are each independently selected from the group consisting of:  
 hydrogen,  
 halogen,  
 alkenyl,  
 amino,  
 -R<sub>11</sub>,  
 30 -O-R<sub>11</sub>,  
 -S-R<sub>11</sub>, and  
 -N(R<sub>9a</sub>)(R<sub>11</sub>);

R<sub>11</sub> is selected from the group consisting of alkyl, alkoxyalkylenyl, hydroxyalkylenyl, aryl, arylalkylenyl, heteroaryl, heteroarylalkylenyl, heterocyclyl, and heterocyclalkylenyl, each of which is unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl; alkoxy; hydroxy; hydroxyalkyl; aryl; aryloxy; arylalkyleneoxy; heteroaryl; heteroaryloxy; heteroarylalkyleneoxy; halogen; haloalkyl; haloalkoxy; mercapto; nitro; cyano; heterocyclyl; amino; alkylamino; dialkylamino; and, in the case of alkyl, heterocyclyl, and heterocyclalkylenyl, oxo;

R<sub>9a</sub> is selected from the group consisting of hydrogen and C<sub>1-4</sub> alkyl;

R<sub>1</sub> is selected from the group consisting of:







-R<sub>4</sub>,  
 -X-R<sub>4</sub>,  
 -X-Y-R<sub>4</sub>,  
 -X-Y-X-Y-R<sub>4</sub>,  
 -X-R<sub>5</sub>,  
 -N(R<sub>1</sub>')-Q-R<sub>4</sub>,  
 -N(R<sub>1</sub>')-X<sub>1</sub>-Y<sub>1</sub>-R<sub>4</sub>, and  
 -N(R<sub>1</sub>')-X<sub>1</sub>-R<sub>5a</sub>;

X is selected from the group consisting of alkylene, alkenylene, alkynylene, arylene, heteroarylene, and heterocyclylene wherein the alkylene, alkenylene, and alkynylene groups can be optionally interrupted or terminated by arylene, heteroarylene or heterocyclylene and optionally interrupted by one or more -O- groups;


X<sub>1</sub> is C<sub>2-20</sub> alkylene;

Y is selected from the group consisting of:

-O-,  
 -S(O)<sub>0-2</sub>-,  
 -S(O)<sub>2</sub>-N(R<sub>8</sub>)-,  
 -C(R<sub>6</sub>)-,  
 -C(R<sub>6</sub>)-O-,  
 -O-C(R<sub>6</sub>)-,  
 -O-C(O)-O-,  
 -N(R<sub>8</sub>)-Q-,



$\text{Y}_1$  is selected from the group consisting of  $-\text{O}-$ ,  $-\text{S}(\text{O})_{0.2}-$ ,  $-\text{S}(\text{O})_2-\text{N}(\text{R}_8)-$ ,




$-\text{N}(\text{R}_8)-\text{Q}-$ ,  $-\text{C}(\text{R}_6)-\text{N}(\text{R}_8)-$ ,  $-\text{O}-\text{C}(\text{R}_6)-\text{N}(\text{R}_8)-$ , and

$\text{R}_1'$  is selected from the group consisting of hydrogen,  $\text{C}_{1-20}$  alkyl,  
 hydroxy- $\text{C}_{2-20}$  alkylenyl, and alkoxy- $\text{C}_{2-20}$  alkylenyl;


$\text{R}_4$  is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl,  
 arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl, heteroarylalkylenyl,  
 heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl wherein the alkyl, alkenyl,  
 alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl,  
 heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl groups  
 can be unsubstituted or substituted by one or more substituents independently selected

from the group consisting of alkyl; alkoxy; hydroxyalkyl; haloalkyl; haloalkoxy; halogen; nitro; hydroxy; mercapto; cyano; aryl; aryloxy; arylalkyleneoxy; heteroaryl; heteroaryloxy; heteroarylalkyleneoxy; heterocyclyl; amino; alkylamino; dialkylamino; (dialkylamino)alkyleneoxy; and, in the case of alkyl, alkenyl, alkynyl, and heterocyclyl, oxo;

$R_5$  is selected from the group consisting of:



$R_{5a}$  is selected from the group consisting of:



$R_6$  is selected from the group consisting of =O and =S;

$R_7$  is  $C_{2-7}$  alkylene;

$R_8$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl, hydroxy- $C_{1-10}$  alkylenyl,  $C_{1-10}$  alkoxy- $C_{1-10}$  alkylenyl, aryl- $C_{1-10}$  alkylenyl, and heteroaryl- $C_{1-10}$  alkylenyl;

$R_9$  is selected from the group consisting of hydrogen and alkyl;

$R_{10}$  is  $C_{3-8}$  alkylene;

A is selected from the group consisting of  $-\text{CH}_2-$ ,  $-\text{O}-$ ,  $-\text{C(O)}-$ ,  $-\text{S(O)}_{0-2}-$ , and  $-\text{N}(-\text{Q}-\text{R}_4)-$ ;

A' is selected from the group consisting of  $-\text{O}-$ ,  $-\text{S(O)}_{0-2}-$ ,  $-\text{N}(-\text{Q}-\text{R}_4)-$ , and  $-\text{CH}_2-$ ;

Q is selected from the group consisting of a bond,  $-\text{C(R}_6\text{)}-$ ,  $-\text{C(R}_6\text{)}-\text{C(R}_6\text{)}-$ ,  $-\text{S(O)}_2-$ ,  $-\text{C(R}_6\text{)}-\text{N(R}_8\text{)}-\text{W}-$ ,  $-\text{S(O)}_2-\text{N(R}_8\text{)}-$ ,  $-\text{C(R}_6\text{)}-\text{O}-$ ,  $-\text{C(R}_6\text{)}-\text{S}-$ , and  $-\text{C(R}_6\text{)}-\text{N(OR}_9\text{)}-$ ;

V is selected from the group consisting of  $-\text{C(R}_6\text{)}-$ ,  $-\text{O}-\text{C(R}_6\text{)}-$ ,  $-\text{N(R}_8\text{)}-\text{C(R}_6\text{)}-$ , and  $-\text{S(O)}_2-$ ;

W is selected from the group consisting of a bond,  $-\text{C(O)}-$ , and  $-\text{S(O)}_2-$ ; and


a and b are independently integers from 1 to 6 with the proviso that  $a + b$  is  $\leq 7$ ;

or a pharmaceutically acceptable salt thereof.

3. The compound or salt of claim 2 wherein  $G_1$  is selected from the group consisting of  $-C(O)-R'$ ,  $\alpha$ -amino- $C_{2-11}$  acyl, and  $-C(O)-O-R'$ .

5

4. A compound of the Formula III:



III

wherein:

10  $G_2$  is selected from the group consisting of:

$-X_2-C(O)-R'$ ,  
 $\alpha$ -aminoacyl,  
 $\alpha$ -aminoacyl- $\alpha$ -aminoacyl,  
 $-X_2-C(O)-O-R'$ ,  
 15  $-C(O)-N(R'')R'$ , and  
 $-S(O)_2-R'$ ;

$X_2$  is selected from the group consisting of a bond;  $-CH_2-O-$ ;  $-CH(CH_3)-O-$ ;  $-C(CH_3)_2-O-$ ; and, in the case of  $-X_2-C(O)-O-R'$ ,  $-CH_2-NH-$ ;

20  $R'$  and  $R''$  are independently selected from the group consisting of  $C_{1-10}$  alkyl,  $C_{3-7}$  cycloalkyl, phenyl, and benzyl, each of which may be unsubstituted or substituted by one or more substituents independently selected from the group consisting of halogen, hydroxy, nitro, cyano, carboxy,  $C_{1-6}$  alkyl,  $C_{1-4}$  alkoxy, aryl, heteroaryl, aryl- $C_{1-4}$  alkylenyl, heteroaryl- $C_{1-4}$  alkylenyl, halo- $C_{1-4}$  alkylenyl, halo- $C_{1-4}$  alkoxy,  $-O-C(O)-CH_3$ ,  $-C(O)-O-CH_3$ ,  $-C(O)-NH_2$ ,  $-O-CH_2-C(O)-NH_2$ ,  $-NH_2$ , and  $-S(O)_2-NH_2$ , with the proviso  
 25 that  $R''$  can also be hydrogen;

$\alpha$ -aminoacyl is an  $\alpha$ -aminoacyl group derived from an amino acid selected from the group consisting of racemic, D-, and L-amino acids;

$R_A$  and  $R_B$  are each independently selected from the group consisting of:  
 hydrogen,



halogen,  
 alkenyl,  
 amino,  
 -R<sub>11</sub>,  
 -O-R<sub>11</sub>,  
 -S-R<sub>11</sub>, and  
 -N(R<sub>9a</sub>)(R<sub>11</sub>);

5

10

15

R<sub>11</sub> is selected from the group consisting of alkyl, alkoxyalkylenyl, hydroxyalkylenyl, aryl, arylalkylenyl, heteroaryl, heteroarylalkylenyl, heterocyclyl, and heterocyclylalkylenyl, each of which is unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl; alkoxy; hydroxy; hydroxyalkyl; aryl; aryloxy; arylalkyleneoxy; heteroaryl; heteroaryloxy; heteroarylalkyleneoxy; halogen; haloalkyl; haloalkoxy; mercapto; nitro; cyano; heterocyclyl; amino; alkylamino; dialkylamino; and, in the case of alkyl, heterocyclyl, and heterocyclylalkylenyl, oxo;

R<sub>9a</sub> is selected from the group consisting of hydrogen and C<sub>1-4</sub> alkyl;

R<sub>1</sub> is selected from the group consisting of:

20

25

-R<sub>4</sub>,  
 -X-R<sub>4</sub>,  
 -X-Y-R<sub>4</sub>,  
 -X-Y-X-Y-R<sub>4</sub>,  
 -X-R<sub>5</sub>,  
 -N(R<sub>1</sub>')-Q-R<sub>4</sub>,  
 -N(R<sub>1</sub>')-X<sub>1</sub>-Y<sub>1</sub>-R<sub>4</sub>, and  
 -N(R<sub>1</sub>')-X<sub>1</sub>-R<sub>5a</sub>;

30






X is selected from the group consisting of alkylene, alkenylene, alkynylene, arylene, heteroarylene, and heterocyclylene wherein the alkylene, alkenylene, and alkynylene groups can be optionally interrupted or terminated by arylene, heteroarylene or heterocyclylene and optionally interrupted by one or more -O- groups;

X<sub>1</sub> is C<sub>2-20</sub> alkylene;


Y is selected from the group consisting of:

-O-,

-S(O)<sub>0-2</sub>-,  
 -S(O)<sub>2</sub>-N(R<sub>8</sub>)-,  
 -C(R<sub>6</sub>)-,  
 -C(R<sub>6</sub>)-O-,  
 5 -O-C(R<sub>6</sub>)-,  
 -O-C(O)-O-,  
 -N(R<sub>8</sub>)-Q-,  
 -C(R<sub>6</sub>)-N(R<sub>8</sub>)-,  
 -O-C(R<sub>6</sub>)-N(R<sub>8</sub>)-,  
 10 -C(R<sub>6</sub>)-N(OR<sub>9</sub>)-,  
 -O-N(R<sub>8</sub>)-Q-,  
 -O-N=C(R<sub>4</sub>)-,  
 -C(=N-O-R<sub>8</sub>)-,  
 -CH(-N(-O-R<sub>8</sub>)-Q-R<sub>4</sub>)-,

15 ,  
,  
,  
, and  
;

20 Y<sub>1</sub> is selected from the group consisting of -O-, -S(O)<sub>0-2</sub>-, -S(O)<sub>2</sub>-N(R<sub>8</sub>)-,


-N(R<sub>8</sub>)-Q-, -C(R<sub>6</sub>)-N(R<sub>8</sub>)-, -O-C(R<sub>6</sub>)-N(R<sub>8</sub>)-, and ;

R<sub>1</sub>' is selected from the group consisting of hydrogen, C<sub>1-20</sub> alkyl,


hydroxy-C<sub>2-20</sub> alkylenyl, and alkoxy-C<sub>2-20</sub> alkylenyl;

R<sub>4</sub> is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl, heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl wherein the alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl, heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl groups can be unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl; alkoxy; hydroxyalkyl; haloalkyl; haloalkoxy; halogen; nitro; hydroxy; mercapto; cyano; aryl; aryloxy; arylalkyleneoxy; heteroaryl; heteroaryloxy; heteroarylalkyleneoxy; heterocyclyl; amino; alkylamino; dialkylamino; (dialkylamino)alkyleneoxy; and, in the case of alkyl, alkenyl, alkynyl, and heterocyclyl, oxo;

R<sub>5</sub> is selected from the group consisting of:



R<sub>5a</sub> is selected from the group consisting of:



R<sub>6</sub> is selected from the group consisting of =O and =S;

R<sub>7</sub> is C<sub>2-7</sub> alkylene;

R<sub>8</sub> is selected from the group consisting of hydrogen, C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, hydroxy-C<sub>1-10</sub> alkylenyl, C<sub>1-10</sub> alkoxy-C<sub>1-10</sub> alkylenyl, aryl-C<sub>1-10</sub> alkylenyl, and heteroaryl-C<sub>1-10</sub> alkylenyl;

R<sub>9</sub> is selected from the group consisting of hydrogen and alkyl;

R<sub>10</sub> is C<sub>3-8</sub> alkylene;

A is selected from the group consisting of -CH<sub>2</sub>-, -O-, -C(O)-, -S(O)<sub>0-2</sub>-, and -N(-Q-R<sub>4</sub>)-;

A' is selected from the group consisting of -O-, -S(O)<sub>0-2</sub>-, -N(-Q-R<sub>4</sub>)-, and -CH<sub>2</sub>-;

Q is selected from the group consisting of a bond, -C(R<sub>6</sub>)-, -C(R<sub>6</sub>)-C(R<sub>6</sub>)-,  
-S(O)<sub>2</sub>-, -C(R<sub>6</sub>)-N(R<sub>8</sub>)-W-, -S(O)<sub>2</sub>-N(R<sub>8</sub>)-, -C(R<sub>6</sub>)-O-, -C(R<sub>6</sub>)-S-, and -C(R<sub>6</sub>)-N(OR<sub>9</sub>)-;

V is selected from the group consisting of -C(R<sub>6</sub>)-, -O-C(R<sub>6</sub>)-, -N(R<sub>8</sub>)-C(R<sub>6</sub>)-, and  
5 -S(O)<sub>2</sub>-;

W is selected from the group consisting of a bond, -C(O)-, and -S(O)<sub>2</sub>-; and

a and b are independently integers from 1 to 6 with the proviso that a + b is ≤ 7;  
or a pharmaceutically acceptable salt thereof.

10 5. The compound or salt of claim 4 wherein G<sub>2</sub> is selected from the group consisting  
of α-amino-C<sub>2-5</sub> alkanoyl, C<sub>2-6</sub> alkanoyl, C<sub>1-6</sub> alkoxy carbonyl, and C<sub>1-6</sub> alkyl carbamoyl.

6. The compound or salt of any one of claims 1 through 5 wherein R<sub>A</sub> and R<sub>B</sub> are  
independently selected from the group consisting of hydrogen, -R<sub>11</sub>, -O-R<sub>11</sub>, and -NHR<sub>11</sub>,  
15 wherein R<sub>11</sub> is alkyl, alkoxyalkylenyl, or hydroxyalkylenyl.

7. The compound or salt of claim 6 wherein R<sub>A</sub> is selected from the group consisting  
of hydrogen and C<sub>1-5</sub> alkyl, and R<sub>B</sub> is selected from the group consisting of C<sub>1-5</sub> alkyl,  
-O-C<sub>1-4</sub> alkyl, and -NH-C<sub>1-4</sub> alkyl.

20

8. The compound or salt of claim 7 wherein R<sub>A</sub> is hydrogen.

9. The compound or salt of claim 8 wherein R<sub>B</sub> is C<sub>1-5</sub> alkyl.

25 10. The compound or salt of claim 7 wherein R<sub>A</sub> and R<sub>B</sub> are each methyl.


11. The compound or salt of any one of claims 1 through 10 wherein R<sub>1</sub> is selected  
from the group consisting of:

30 -R<sub>4</sub>,  
-X-R<sub>4</sub>,  
-X-Y-R<sub>4</sub>,  
-X-Y-X-Y-R<sub>4</sub>, and

-X-R<sub>5</sub>.

12. The compound or salt of claim 11 wherein R<sub>1</sub> is -R<sub>4</sub> or -X-R<sub>4</sub>.

5 13. The compound or salt of claim 12 wherein -X- is



14. The compound or salt of claim 12 wherein R<sub>1</sub> is selected from the group consisting of aryl-C<sub>1-4</sub> alkylenyl and heteroaryl-C<sub>1-4</sub> alkylenyl, wherein the aryl or heteroaryl group is  
 10 unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl, alkoxy, hydroxyalkyl, haloalkyl, haloalkoxy, halogen, nitro, hydroxy, mercapto, cyano, aryl, aryloxy, arylalkyleneoxy, heteroaryl, heteroaryloxy, heteroarylalkyleneoxy, heterocyclyl, amino, alkylamino, dialkylamino, and (dialkylamino)alkyleneoxy.

15 15. The compound or salt of claim 14 wherein R<sub>1</sub> is benzyl, which is unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl, alkoxy, haloalkyl, haloalkoxy, and halogen.

20 16. The compound or salt of claim 15 wherein R<sub>1</sub> is benzyl or 4-fluorobenzyl.


17. The compound or salt of claim 12 wherein R<sub>1</sub> is tetrahydro-2*H*-pyran-4-ylmethyl.

18. The compound or salt of claim 12 wherein R<sub>1</sub> is pyridin-3-ylmethyl, isoxazol-5-ylmethyl, isoxazol-3-ylmethyl, [5-(4-fluorophenyl)isoxazol-3-yl]methyl, or [3-(4-fluorophenyl)isoxazol-5-yl]methyl.  
 25

19. The compound or salt of claim 11 wherein R<sub>1</sub> is -X-Y-R<sub>4</sub>.

30 20. The compound or salt of claim 19 wherein R<sub>1</sub> is -C<sub>2-5</sub>alkylenyl-S(O)<sub>2</sub>-C<sub>1-3</sub>alkyl.

21. The compound or salt of claim 19 wherein R<sub>1</sub> is



22. The compound or salt of claim 19 wherein R<sub>1</sub> is -C<sub>2-5</sub>alkylenyl-NH-Q-R<sub>4</sub>.

5

23. The compound or salt of claim 21 or 22 wherein Q is -C(O)-, S(O)<sub>2</sub>-, or -C(O)-NH- and R<sub>4</sub> is C<sub>1-6</sub>alkyl.

24. A pharmaceutical composition comprising a therapeutically effective amount of a  
10 compound or salt of any one of claims 1 through 23 and a pharmaceutically acceptable carrier.

25. A method of inducing cytokine biosynthesis in an animal comprising administering  
an effective amount of a compound or salt of any one claims 1 through 23 or the  
15 pharmaceutical composition of claim 24 to the animal.

26. A method of selectively inducing the biosynthesis of IFN- $\alpha$  in an animal  
comprising administering an effective amount of a compound or salt of any one claims 1  
through 23 or the pharmaceutical composition of claim 24 to the animal.

20

27. A method of treating a viral disease in an animal comprising administering a  
therapeutically effective amount of a compound or salt of any one of claims 1 through 23  
or the pharmaceutical composition of claim 24 to the animal.

28. A method of treating a viral disease in an animal comprising administering a  
therapeutically effective amount of a compound or salt of any one of claims 1 through 23  
or the pharmaceutical composition of claim 24 to the animal; and selectively inducing the  
25 biosynthesis of IFN- $\alpha$  in the animal.

29. A method of treating a neoplastic disease in an animal comprising administering a therapeutically effective amount of a compound or salt of any one of claims 1 through 23 or the pharmaceutical composition of claim 24 to the animal.
- 5 30. A method of treating a neoplastic disease in an animal comprising administering a therapeutically effective amount of a compound or salt of any one of claims 1 through 23 or the pharmaceutical composition of claim 24 to the animal; and selectively inducing the biosynthesis of IFN- $\alpha$  in the animal.

# INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2006/034427

## A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D471/04 A61K31/437 A61P37/02

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FRANKOWSKI, A.: "Synthesis of imidazo[4,5-c]pyridine and imidazo[4,5-d][1,2]diazepine systems and their ribonucleosides" TETRAHEDRON, 42(5), 1511-28 CODEN: TETRAB; ISSN: 0040-4020, 1986, XP002413466 compound 28 of page 1515 page 1522, line 41 - page 1523, line 3	2,3,6, 11-16
X	DATABASE CHEMCATS CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US.; XP002413928 Order Number: PHAR090799 & "AMBINTER STOCK SCREENING COLLECTION" 3 July 2005 (2005-07-03), AMBINTER, PARIS, FRANCE	2,3,6, 11,12



Further documents are listed in the continuation of Box C.



See patent family annex.

### \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

10 January 2007

Date of mailing of the international search report

24/01/2007

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

MATES VALDIVIELSO, J



## INTERNATIONAL SEARCH REPORT

International application No

PCT/US2006/034427

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	YUTILOV, YU.M. & SVERTILOVA, I.A.: "Synthesis and some reactions of 4-nitro derivatives of imidazo[4,5-c]pyridin-2-ones" CHEMISTRY OF HETEROCYCLIC COMPOUNDS, vol. 22, no. 1, 1986, pages 80-85, XP008073281 New York compounds VIIA, VIIB page 84, paragraph 5-8	1,6,11, 12
A	WO 2005/051317 A2 (3M INNOVATIVE PROPERTIES CO [US]; KREPSKI LARRY R [US]; DELLARIA JOSEP) 9 June 2005 (2005-06-09) abstract claim 1	1,2,4, 24-30
P,X	WO 2006/044573 A2 (GEN ATOMICS [US]; YUAN CHONG-SHENG [US]) 27 April 2006 (2006-04-27) paragraph [0124] claims 4,24-27	1-30

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US2006/034427

### Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  

Although claims 25-30 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

### Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2006/034427

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2005051317	A2	09-06-2005	AR 046781 A1 21-12-2005
			AU 2004293078 A1 09-06-2005
			CA 2547020 A1 09-06-2005
			EP 1687307 A2 09-08-2006
			MX PA06005910 A 23-08-2006
WO 2006044573	A2	27-04-2006	NONE